

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C12N 7/00, A61K 39/21 G01N 33/569, C12N 15/00 A61K 39/395, C12Q 1/68, 1/70	A1	(11) International Publication Number: WO 89/12094 (43) International Publication Date: 14 December 1989 (14.12.89)
(21) International Application Number: PCT/EP89/00643 (22) International Filing Date: 8 June 1989 (08.06.89) (30) Priority data: 9 June 1988 (09.06.88) EP 88109200.1 <i>(34) Country for which the regional or international application was filed:</i> DE (71) Applicant (for all designated States except US): INNOGEN-ETICS N.V. [BE/BE]; Industriepark Zwijnaarde 7, Box 8, B-9710 Gent (BE). (72) Inventors; and (75) Inventors/Applicants (for US only) : DE LEYS, Robert [BE/BE]; Diepekanten 20, B-1860 Grimbergen (BE). VAN-DEBORGHT, Bart [BE/BE]; Kraaibossen 24, B-2440 Geel (BE). SAMAN, Eric [BE/BE]; Nachtegalenlaan 12, B-2680 S.-Niklaas (BE). VAN HEUVERSWYN, Hugo [BE/BE]; Colmanstraat 62, B-9280 Laarne (BE).		(74) Agents: PAGENBERG, Jochen et al.; Galileiplatz 1, D-8000 München 80 (DE). (81) Designated States: AU, BR, DK, JP, US. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HIV-3 RETROVIRUS AND ITS USE (57) Abstract <p>Described is a new retrovirus designated HIV-3 and deposited in the European Collection of Animal Cell Cultures (ECACC) under V88060301. Further described are antigens obtained from the virus, particularly proteins p12, p16, p26 and glycoproteins gp41 and gp120 to be used in the diagnosis of ARC or AIDS caused by HIV-3. Immunogenic compositions to be used as vaccines contain an envelope glycoprotein of HIV-3 such as gp41 or gp120.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Fasso	HU	Hungary	NO	Norway
BG	Bulgaria	IT	Italy	RO	Romania
BJ	Benin	JP	Japan	SD	Sudan
BR	Brazil	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

HIV-3 retrovirus and its use

-1-

1 Substantial progress has been made in our understanding of
the acquired immunodeficiency syndrome or AIDS. The
principal causative agent has been demonstrated to be a
5 non-transforming retrovirus with a tropism for T4
helper/inducer lymphocytes (1,2) and it has been estimated
that millions of people world-wide have already been
infected. Infection with this virus leads, at least in a
significant percentage of cases, to a progressive depletion
10 of the T4 lymphocyte population with a concomittant
increasing susceptibility to the opportunistic infections
which are characteristic of the disease.
Epidemiological studies indicate that human immunodeficiency
virus type 1 (HIV-1), the etiological agent responsible for
15 the majority of AIDS cases and which is currently the most
widely disseminated HIV, probably had its origins in Central
Africa (3). The discovery of this virus did not necessarily
imply the existence of other types of human immunodeficiency
viruses. Nevertheless, a second group of human
20 immunodeficiency-associated retroviruses, human
immunodeficiency virus type 2 (HIV-2), was identified in West
Africa (4,5). An HIV-2 virus is disclosed in EP-A-0 239 425.
An HIV-1 virus is disclosed in WO 86/02383. Other similar,
but not identical, retroviruses have also been isolated from
25 simian sources (simian immunodeficiency virus, SIV) such as
African green monkeys (6,7) and macaques (8,9). The simian
isolates have been shown to be genetically more closely
related to HIV-2 than HIV-1 but are nevertheless distinct
(10).
30 One characteristic of human immunodeficiency viruses which
complicates their comparison is their genetic variability;
genetic variants arise spontaneously and with high
frequency. A comparison of various HIV-1 isolates revealed
that some regions of the genome are highly variable while
others are reasonably well conserved (11-16). Similar
35 polymorphisms have also been observed for HIV-2 (17). The

-2-

1 regions with the greatest genetic stability are presumably
those regions coding for the regions of viral proteins which
are structurally or enzymatically essential. The viral
5 genes with the greatest overall genetic stability are the
gag and pol genes, while some regions of the env gene and
the genes coding for regulatory proteins such as art, tat,
sor and 3'orf exhibit a high degree of variability. Some of
the major structural features of the gag and pol gene
10 products are apparently shared not only by all of the
variants of a particular HIV type, but have, at least to
some extent, been conserved between virus types. Antiserum
produced against HIV-1 crossreacts with the gag and pol gene
products of HIV-2, albeit with a lower affinity than for the
15 corresponding HIV-1 gene products. However, in spite of the
demonstrable immunological crossreaction, at the nucleic
acid level there is little sequence homology and no
significant hybridization between these two viruses can be
detected except under very low stringency conditions (17).

20 A higher degree of relatedness can be demonstrated between
SIVagm (STLV-III agm, nearly or completely identical to
Human Lymphotropic Virus type 4 (15)) and HIV-2.
Immunological crossreaction is not limited only to the gag
25 and pol gene products but extends to the env gene products
as well. Nevertheless, genomic analysis of SIVagm and HIV-2
showed them to be genetically distinguishable (19). DNA
probes specific for HIV-2, although able to hybridize to
SIVagm sequences, hybridize preferentially to HIV-2 (18).

30 We now report the isolation and characterization of a novel
human immunodeficiency virus from a Camerounian woman and
her partner. Geographically, this virus comes from a region
in Africa located between West Africa where HIV-2 is
35 endemic, and East-Central Africa where HIV-1 is endemic.
This isolate is shown immunologically to be antigenically

-3-

1 more closely related to HIV-1 than is HIV-2, yet an analysis
of partial cleavage products obtained by chemical cleavage
of the gag and pol gene products demonstrate that this
5 isolate is neither HIV-1 nor HIV-2. This novel isolate
could represent an evolutionary link between HIV-1 and
HIV-2. This novel virus will be referred to as HIV-3
hereinafter.

10 Accordingly, the invention relates to an HIV-3 retrovirus or
variants of this virus having the essential morphological
and immunological properties of the retrovirus deposited in
the European Collection of Animal Cell Cultures (ECACC)
under V 88060301.

15 A virus isolation was performed from blood from an
asymptomatic Camerounian woman who is the partner of an
HIV-seropositive man with generalized lymphadenopathy.
Serum from the woman was moderately positive (ratio
O.D./cut-off of 4.5) in the enzyme-linked immunosorbent
20 assay (EIA, Organon Teknika) and had a low titer (1/40) in
the immunofluorescent antibody assay for HIV-1 but gave
ambiguous results in the HIV-1 Western blot assay with clear
bands at p33, P53/55 and p64 but very weak bands at p24,
gp41 and gp120. The virus was isolated by co-cultivation of
25 the woman's lymphocytes with PHA-stimulated lymphocytes from
healthy uninfected donors in a medium consisting of RPMI
1640 buffered with 20 mM HEPES (hydroxyethylpiperazine
ethanesulfonate) and supplemented with 15 % fetal calf
serum, 5 g/ml hydrocortisone, 75 u/ml interleukin-2 (IL-2)
30 and 2 g/ml polybrene.

After 52 days in culture, virus was detected in the culture
as judged by the presence of syncytia and on the basis of
positive immunofluorescence observed when a laboratory
reference anti-HIV antiserum was incubated with
35 acetone-fixed cells from the culture. The presence of
reverse transcriptase was also detected in the culture

-4-

1 supernatant (10^4 cpm/ml, 27 X background). Cell-free
culture supernatant was used to passage the virus on fresh
lymphocytes. After 15 days, CPE was again observed and
5 reverse transcriptase detected in the supernatant. The
virus was further propagated in PHA-stimulated lymphocytes
from healthy blood donors and was transferred to continuous
cell lines of leukemic origin. Virus-containing supernatant
was tested in parallel with culture supernatants known to
10 contain HIV-1 in the differential antigen capturing test
which is described in detail below. The results of
this comparison indicated that the new isolate was not
HIV-1.

15 The new virus was then characterized with respect to its
protein antigens and nucleic acids. The cell lines used for
propagating the virus can be, depending on the case, lines
of the CEM, HUT, Molt-4, or MT4 type, or any other
immortalized cell line which bears the T4 receptor on its
20 cell surface.

A preferred cell line for the continuous propagation of
HIV-3 is Molt-4. Molt-4 cells infected with HIV-3 were
deposited with the ECACC on June 3, 1988 under number V
88060301. Establishment of a chronically-infected cell line
25 can, for example, be carried out as follows:

Molt-4 cells (10^6 /ml) and preferably Molt-4 clone 8 cells
(obtained from N. Yamamoto, Yamaguchi, Japan) are cocultured
with infected human lymphocytes (10^6 /ml) in RPMI 1640
30 culture medium buffered with 20 mM HEPES and containing 10 %
fetal calf serum. Within one to two weeks, a cytopathic
effect is observed in the culture which is followed by cell
death. A fraction of the cells in the culture survive the
infection and produce virus continuously. With continued
35 culturing, these cells increase in number and can be
passaged. Supernatants from these cells can be used a a

-5-

1 source of virus.

Furthermore, the invention relates to a purified retrovirus having the essential morphological and immunological
5 properties described below. In many cases, the unique characteristics of HIV-3 can best be appreciated by comparison with the same type of characteristics relating to the other human immunodeficiency viruses, HIV-1 and HIV-2.

10 Brief description of the drawings

In Figures 1 to 16 the designations HIV-3 (ANT 70) and HIV-3 (ANT 70 NA) refer to two strains of a new HIV-3 virus
15 isolated from a Camerounian woman and her partner from which HIV-3 (ANT 70) has been deposited under ECACC V88060301.

Figure 1 shows a procedure for preparing cleavage maps of viral proteins.

20 Figure 2 shows differential antigen capturing on virus-containing culture supernatants.

Differential antigen capturing is performed as described
25 hereinafter. The solid line represents the results obtained using a broad-spectrum anti-HIV-1 IgG while the broken line depicts the results obtained using an IgG which was rather specific for HIV-1. The titrations shown in panels A-E are typical for HIV-1. Panel F shows the result obtained with
30 HIV-3 (ANT 70) containing supernatant.

Figure 3 shows differential antigen capturing on HIV-1 and HIV-3 (ANT 70 NA) supernatants.

35 Differential antigen capturing was performed as described hereinafter. The solid line depicts the results obtained on

-6-

1 plates coated with the broad spectrum anti-HIV IgG while the
broken line represents the results obtained on plates coated
5 with IgG which shows less crossreactivity with HIV types
other than HIV-1.

Figure 4 shows the reactivity of anti-HIV sera on HIV-1 and
HIV-2 Western Blot strips.

10 The reactivities of 3 different sera on HIV-1 and HIV-2
Western blot strips are shown. Sera: 1. anti-HIV-1, 2.
anti-HIV-3 (ANT 70), 3. anti-HIV-2 (isolate 53). The
molecular weights indicated are those given by the
manufacturer (Dupont Biotech).

15 Figure 5 relates to the comparison of gag and pol proteins
of several HIV-1 isolates, HIV-2rod and HIV-3 (ANT 70).

20 Proteins were separated electrophoretically and blotted as
described later. The blot was incubated with a
broad-spectrum anti- HIV antiserum followed by (anti-human
IgG)/alkaline phosphatase- labeled conjugate to visualize
the proteins.

25 A. HIV-2rod, B. an HIV-1 laboratory isolate, C. HIV-3 (ANT
70), D. an HIV-1 laboratory isolate, E. HIV-1 (SF4).

Figure 6 shows a comparison of HIV-3 (ANT 70) and HIV-3 (ANT
70 NA) proteins.

30 Proteins were separated electrophoretically and blotted as
described later. The blot was incubated with the BSR
antiserum followed by (alkaline phosphatase)/anti-human IgG
conjugate to visualize the proteins. Lane 1: HIV-3 (ANT 70
NA), lane 2: HIV-3 (ANT 70), lane 3: HIV-1 (SF4). The
35 apparent intensity difference between lanes 1 and 2 is
caused by the difference in the amount of material loaded.

-7-

1 Figure 7 relates to the ability of various human anti-HIV-1 sera to capture viral antigens.

5 A number of human sera were diluted 1:1000 and coated directly on microwell plates. Detergent-treated culture supernatants containing HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod or HIV-2 (isolate 53) were incubated and the bound
10 antigen was detected using a broadspectrum (anti-HIV)/horseradish peroxidase conjugate. Sera 1-7 were of African origin while sera 8-11 were from Europeans. The greater ability of African sera to capture non-HIV-1 antigen can, in part, be explained by their higher anti-p24 titers (data not
15 shown).

Figure 8 shows the effect of coating IgG dilution on the binding of HIV isolates.

20 Succesive 2-fold dilutions were made of four different sera, beginning at a dilution of 1:1000 and were used to coat microwell plates. Detergent-treated supernatants of HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod and HIV-2 (isolate 53) were diluted to give approximately the same optical density on plates coated with the antiserum shown in panel B at a
25 dilution of 1:1000. Bound antigen was detected using the broad-spectrum (anti-HIV IgG)/horseradish peroxidase conjugate.

30 Figure 9 shows antigen capturing of virus isolates using human polyclonal and mouse anti-HIV-1 monoclonal antibodies.

Wells were coated and incubated as described in the text. The IgGs used are as follows:

35 1. human polyclonal anti-HIV IgG , 2. MAb CLB 59, 3. MAb CLB

-8-

1
21, 4. MAb CLB 64, 5. MAb CLB 14, 6. MAb CLB 16, 7. MAb CLB
47, 8. MAb CLB 13.6 (anti-p18), 9. MAb CLB 19.7, 10. MAb CLB
13.4 (anti-p18).

5
Figure 10 is a comparison of the reactivity of human
anti-HIV antisera to different HIV types.

10
Lysates of HIV-1 (SF4), HIV-3 (ANT 70), HIV-2rod and HIV-2
(isolate 53) were separated electrophoretically on
SDS-polyacrylamide gels, blotted onto nitrocellulose, and
incubated with a high titer anti-HIV-1 antiserum (panel A),
a lower titer anti-HIV-1 antiserum (panel B), serum from the
15 woman from whom HIV-3 (ANT 70) was isolated (panel C), her
partner from which HIV-3 (ANT 70 NA) was isolated (panel D)
and anti-HIV-2 antiserum from the person from whom HIV-2
(isolate 53) was isolated (panel E).

20
Figure 11 shows titrations of anti-HIV sera by enzyme
immunoassay.

25
Microwell plates were coated with lysates of HIV-1 (SF4),
HIV-3 (ANT 70) and HIV-2 (isolate 53). Serum from an
HIV-1-infected European (left panel), antiserum to HIV-3
(ANT 70 NA) (center panel) and antiserum to HIV-2 (isolate
53) (right panel) were titrated in 2-fold dilutions
beginning at a dilution of 1:100 on all three coated plates.

30
Figure 12 shows the positions of methionine and tryptophan
residues in viral gag and pol gene products.

35
Amino acid positions for the p17 gag proteins are given
starting from the first methionine in the coding sequence.
Positions for the p24 gag protein are given starting at the
p17/p24 proteolytic cleavage site. Positions for the pol
gene are shown after alignment with the highly conserved

-9-

1 tryptophan doublet in the HIV-1 sequence at positions 556
and 557. The positions of a conserved protease sequence,
the protease/reverse transcriptase cleavage site and the
5 reverse transcriptase/endonuclease cleavage site are
indicated. In this case, the terms p24 and p17 are used in
the genetic sense to refer to the largest and second largest
viral core proteins respectively. The term "HIV-2 (LAV-2)" is
a synonymum for HIV-2 rod.

10 Figure 13 is a comparison of partial cleavage products of
gag and pol gene products of HIV-1 (SF4) [HIV-1 in the
figure], HIV-3 (ANT 70) [isolate 70 in the figure], HIV-2rod
[HIV-2 (LAV-2) in the figure] and HIV-2 (isolate 53)
15 [isolate 53 in the figure]. The terms p24 and p17 are used
in the genetic sense to indicate the largest and second
largest viral core proteins, respectively.

Figure 14 shows hybridization of cDNA probes to viral RNA.

20 Viral RNA from HIV-1 (SF4), HIV-2rod, and HIV-3 (ANT 70)
were spotted onto a membrane filter as described in
Materials and Methods. The filters were hybridized under
either nonstringent (A) or stringent conditions and
25 autoradiographed.

1. Morphology

30 Electron microscopy of HIV-3-infected MT4 cells revealed the
presence of extracellular virus particles having a diameter

35

-10 -

1 of approximately 120nm and consisting of an outer envelope
which surrounds an inner elongated core which has a diameter
of approximately 20 to 40 nm and which appears in some thin
5 sections to be slightly cone-shaped in contrast to the more
or less cylindrical appearance of the HIV-1 core.
Nevertheless, HIV-3 is morphologically very similar to HIV-1
and HIV-2 but is readily distinguished from other human
retroviruses such as HTLV-I and HTLV-II.

10 2. Protein and glycoprotein antigens.

The virus present in the culture supernatant of
HIV-3-infected Molt-4 cells was concentrated by
15 precipitation with polyethyleneglycol (average molecular
weight 6000) followed by centrifugation. The resulting
pellet was resuspended in phosphate buffered saline, layered
on top of a 20 % sucrose cushion and pelleted at 100,000 g
for 1.5 hours. The pelleted virus was then dissociated in
20 62.5 mM Tris, pH 6.7, containing 2 % 2-mercaptoethanol, 1 %
sodium dodecyl sulfate and 10 % glycerol and the principle
viral antigens were separated by electrophoresis on a
polyacrylamide gel (12,5 %) under denaturing conditions.
Molecular weight markers were included on the same gel so as
25 to provide a basis for estimating molecular weights. Once
separated, the proteins were electrophoretically transferred
to nitrocellulose paper (Western blot) which was then
incubated with an antiserum derived from a person infected
with an HIV. In the initial experiments, a high titer
30 antiserum was used from an individual who was infected with
HIV-1 and which had been previously shown to crossreact with
HIV-2 gag- and pol-derived proteins. In this manner, the
molecular weights of the HIV-3 gag and pol gene products
could be compared with those of HIV-1 and HIV-2.

35 The apparent molecular weights observed for the HIV-3
proteins are close to those observed for both HIV-1 and

-11-

1 HIV-2. Nevertheless, small yet reproducible molecular weight differences between HIV-3, and HIV-1 and HIV-2 proteins are also evident.

5 The protein blots revealed that HIV-3, like HIV-1 and HIV-2, possesses three core proteins. In the case of HIV-3, these proteins were found to have molecular weights of approximately 12,000, 16,500 and 25,000 respectively. By convention, proteins are frequently referred to by a "p" for protein or "gp" for glycoprotein, followed by a number
10 which, when multiplied by 1,000, gives the approximate molecular weight of the polypeptide. The three major core proteins of HIV-3 will be referred to hereafter as p12, p16, and p25 respectively.

15 The molecular weight values as determined are expected to be correct to within 10 % of the true values. Nevertheless, much confusion exists with regards to molecular weight values of proteins since the construction of the electrophoresis apparatus used and the source of the buffer components varies from laboratory to laboratory. It is
20 therefore necessary when comparing the apparent molecular weights of the protein antigens of HIV-3 with respect to those of HIV-1 or HIV-2, to subject all samples to electrophoresis on the same gel. Such a gel can, for example, be seen in Fig. 5. In particular, it is
25 evident that while, in the case of the major core protein, the molecular weight values of the homologous proteins of the three HIVs are very close, the protein derived from HIV-1 is the smallest. The major core protein of HIV-2 is somewhat larger than that of HIV-1, as has been previously
30 reported. The homologous protein from HIV-3 is slightly larger than the major core protein of HIV-2. The calculated molecular weights of these proteins are given in Table 1.

35

-12-

Table 1 - Comparison of molecular weights of gag and pol gene products.

	<u>gag</u>				<u>pol</u>		<u>env</u>	
					ENDO	REVERSE TRANSCRIP- TASE**	TRANS MEMBRANE PROTEIN	OUTER MEMBRANE PROTEIN
HIV-1	12 KD	17* KD,	24 KD	31 KD	49	KD/65 KD	gp41	gp120
ANT 70	12 KD	16.5 KD,	24.8 KD	31 KD	48.5	KD/62 KD	gp41	gp120
HIV-2	12 KD	16 KD,	24.3 KD	31 KD	53	KD/69 KD	gp41	gp120

* Some strain to strain variation in molecular weight has been observed for this protein.

** Molecular weights are given for both species of reverse transcriptase.

Similarly, molecular weight differences are also apparent between the three HIVs with respect to the second core protein which has, in most HIV-1 strains, a molecular weight of 18,000. Strain to strain differences in the molecular weight of this protein have, however, been documented in the case of HIV-1, and the molecular weight of this protein may be 17,000 in some isolates. The homologous protein from HIV-2 has a molecular weight of approximately 16,000 while the HIV-3 protein has an intermediate molecular weight of approximately 16,500.

By analogy with HIV-1 and HIV-2, HIV-3 also possesses two

-13 -

1 forms of the virally encoded enzyme reverse transcriptase.
These two species also differ slightly in molecular weight
from the corresponding species in HIV-1 and HIV-2 and are
5 characteristic for HIV-3. These molecular weights are also
summarized in Table 1.

HIV-3 possesses an additional pol gene-derived polypeptide
which is an endonuclease with apparent molecular weight of
31,000 and which does not differ significantly in molecular
10 weight from the homologous proteins from HIV-1 or HIV-2.

When protein blots containing HIV-3 proteins are incubated
with serum obtained from an individual infected with this
virus, two additional proteins can be seen. These proteins
15 are derived from the env gene and are the viral envelope
glycoproteins. The smallest protein, which is the
transmembrane protein, migrates as a broad band with an
apparent molecular weight of between 40,000 and 45,000.
This protein will henceforth be referred to as gp41, with
20 the understanding that the protein exhibits some intrinsic
heterogeneity with respect to its apparent molecular weight
and migration on polyacrylamide gels. The larger protein,
which is the outer membrane protein, is similarly somewhat
diffuse on polyacrylamide gels and has a molecular weight of
25 approximately 120,000. This protein will henceforth be
referred to as gp120. It should be noted that the apparent
heterogeneous migration of these two species on
polyacrylamide gel is not due to heterogeneity in the
polypeptide chain but rather in posttranslational
30 glycosylation. In particular, the gp120 is heavily
glycosylated and the apparent molecular weight which one
observes is to some degree influenced by the cell line used
to produce the virus.

35 In addition to the Western blot, viral protein antigens can
also be visualized by radioimmunoprecipitation assay (RIPA).

-14-

1 For this purpose, viral proteins can be radioactively
labeled metabolically in vivo by culturing HIV-3-infected
cells in the presence of 35S-cysteine and 35S-methionine
5 (200 Ci/ml) in RPMI 1640 medium devoid of these two amino
acids and supplemented with dialyzed fetal calf serum.
After 16 hours, the labeled virus is harvested from the
culture supernatant by centrifugation over a 20 % sucrose
cushion at 100,000 g for 1,5 hours. The resulting pelleted
10 virus is then resuspended in RIPA buffer (20 mM
triethanolamine, pH 8.0, 0.5 M NaCl, 0.5 % Nonidet P40, 0.1
% sodium deoxycholate, and 1 mM
phenylmethylsulfonylfluoride).
Alternatively, the virus may be radioactively labeled with
15 125 I using chloramine T by the technique familiar to
persons versed in the art. In this case, virus is purified
from the supernatant of infected cells by pelleting the
virus through a cushion of 20 % sucrose, resuspending the
virus in phosphate buffered saline and banding the virus by
20 ultracentrifugation on a 20 to 50 % sucrose gradient at
60,000 g for 12 hours. The banded virus can be located in
the fractionated gradient either by reverse transcriptase
assay or by an antigen capturing assay. The fractions
containing virus are pooled and Triton X-100 is added to a
25 concentration of 0.5 %. The Triton X-100- lysed virus may
then be iodinated.

For immunoprecipitations, 100,000 -200,000 cpm of labeled
viral protein in RIPA buffer is reacted with 5 microliters
30 of a test serum in a volume of 200 microliters for 16 hours
at 4°C. The resulting immune complexes are then bound to
Protein A-Sepharose (Pharmacia), washed extensively, and the
bound proteins eluted with electrophoresis sample buffer
containing 1 % SDS. The antigens are subsequently analyzed
35 by electrophoresis followed by autoradiography.

-15-

1 The protein antigens of HIV-3 can be characterized with
respect to those of HIV-1 and HIV-2 using two different but
related approaches. On one hand, the antigens may be
5 characterized on the basis of their ability to crossreact
with antisera from persons infected with HIV-1 and HIV-2.
On the other hand, antisera from persons infected with
HIV-3, which contain antibodies produced in response to
HIV-3 antigens, can be used to test crossreactivity to HIV-1
10 and HIV-2 proteins. The antigenic relationships between
HIV-3, and HIV-1 and HIV-2 are substantially illustrated in
the examples given below.

The results of these experiments indicate that HIV-3 is only
distantly related to HIV-2 since crossreactivity is only
15 observed with respect to the viral core proteins and pol
gene products. No crossreactivity of the env gene products
was observed when anti-HIV-2 antiserum was incubated with
HIV-3 proteins or when anti-HIV-3 antiserum was incubated
with HIV-2 proteins.

20 In contrast, HIV-3 is more closely related to HIV-1 since
anti- HIV-3 antiserum crossreacts not only with the gag and
pol gene products of HIV-1 but also to some extent with the
gp41 and gp120 env gene products, albeit with a lower
affinity. Anti- HIV-1 antiserum similarly crossreacts with
25 all of the protein antigens of HIV-3, but with a lower
affinity than for the proteins of HIV-1.

In the examples which follow, it is demonstrated that HIV-3
is antigenically substantially different from HIV-1 on the
30 basis of 1.) a different pattern of reactivity with
anti-HIV-1 antiserum than that observed for HIV-1, 2.) a
drastically reduced ability to be recognized by mouse
monoclonal antibodies raised against the HIV-1 p24 and p17
core proteins, and 3.) preferential recognition of HIV-3
35 proteins, including the envelope proteins, over HIV-1
proteins by antisera from HIV-3- infected individuals.

SUBSTITUTE SHEET

-16-

1
5 In spite of the genetic variation characteristic of human immunodeficiency viruses, a test based, for example, on HIV-1 proteins derived from a particular strain will function satisfactorily for detecting antibodies raised in response to other HIV-1 variants. This can, in particular, also be seen in the example in which monoclonal antibodies were tested for their ability to react with antigens derived from HIV-1, HIV-2 and HIV-3 isolates. In this case, the monoclonal antibodies were raised against the core proteins from the HIV-1 IIIB strain, yet react very strongly to proteins derived from HIV-1 strain SF4. In contrast, these same monoclonal antibodies react only weakly or not at all with HIV-3 core proteins. This indicates that the antigenic differences between HIV-1 and HIV-3 are of such a magnitude that immunological assays based on the use of HIV-1 proteins will not be suitable for testing sera from individuals infected with HIV-3.

10
15
20 Finally, in the examples given below, differences have been shown in the number and/or positions of methionine and tryptophan residues in the most highly conserved gag and pol gene products.

25
3. HIV-3 NUCLEIC ACIDS

30 A. HIV-3 viral RNA.

The RNA of HIV-3 when deposited on a Hybond-H (Amersham) filter according to the "dot blot" technique, did not hybridize to HIV-1 DNA under stringent hybridization conditions.

35 By "stringent conditions" or "nonstringent conditions" are

-17-

1 meant the conditions under which the actual hybridization and/or the subsequent wash steps are performed.

5 Dot blot hybridizations were performed by spotting dilutions of viral RNA from HIV-1 strain SF4, HIV-2 rod and HIV-3 strain ANT 70 onto Hybond-H filters.

10 The dilution series for each virus corresponded to viral RNA pelleted from the equivalent of 5, 2.5, 1.25 and 0.62 milliliters of culture supernatant. The RNA was fixed onto the filter by U.V. irradiation for 2 min and subjected to hybridization by bringing the filter into contact with a 32P- labeled DNA probe. The probe chosen was derived from the HIV-1 sequence spanning nucleotides 487-4652 (Sac I - Eco R1) and includes a portion of the 5' long terminal repeat, the entire gag region and most of the pol gene, 15 subcloned in the vector pUC 13. Hybridization of the 32P-labeled probe with the filter was carried out under stringent conditions in 3 X SSC, 0.5 % milk powder, 1 % SDS, 10 % dextran sulfate, 50 % formamide (volume/volume) at 20 42°C for 18 hrs (1 X SSC corresponds to 0.15 M NaCl, 0.015 M sodium citrate). The subsequent wash steps were carried out under stringent conditions in 0.1 X SSC and 0.1 % SDS at 65°C (2 - 30 minute washes). The filter was then dried and autoradiographed with enhancing screens at -70°C. Following 25 autoradiography, only spots were visible which corresponded to HIV-1 viral RNA. No hybridization was observed to HIV-2 or either of the two HIV-3 strains. HIV-3 therefore appears to be only distantly related to HIV-1.

30 B. cDNA and subclones of cDNA derived from HIV-3.

35 The conditions under which cDNA corresponding to HIV-3 sequences was synthesized and cloned are described below. HIV-3 (strain ANT 70) from 1 liter of culture was precipitated with polyethylene glycol 6000, redissolved in phosphate buffered saline, and pelleted through a 20 %

-18-

1 sucrose cushion. The resulting virus pellet was dissolved
in 6 M guanidinium chloride in 20 mM dithiothreitol and 0.5
% Nonidet P-40. CsCl was added to a concentration of 2 molar
5 and the solution containing disrupted virus was layered onto
a 1.2 milliliter cushion of 5.7 M CsCl containing 0.1 EDTA.
Viral RNA was pelleted by centrifuging for 20 hrs. at 25,000
rpm in a Beckman SW28 rotor at 15°C. The pelleted RNA was
redissolved, extracted with phenol and precipitated with
10 ethanol and 2 M LiCl.

One-fifth of the viral RNA, prepared as described above, was
used to direct the first step in the synthesis of cDNA which
made use of an oligo (dT) primer which served to prime the
synthesis of the first cDNA strand.

15 A commercially available kit supplied by Amersham was used
for preparation of HIV-3 cDNA and made use of an exogenously
added reverse transcriptase to synthesize the first strand.
The synthesis of the second strand was performed using E.
coli DNA polymerase I in the presence of RNase H to digest
20 away the RNA strand of the RNA/DNA hybrid.

Second strand synthesis was performed in the presence of
32p- dCTP to label the cDNA. The resulting cDNA was treated
with T4 DNA polymerase to create blunt ends, the cDNA was
methylated to protect possible internal EcoRI cleavage
25 sites, and was then coupled to EcoRI linkers, also supplied
by Amersham. The EcoRI restriction sites were then cleaved
and the cDNA was sized on a 1.2 % agarose gel. The region
in the gel corresponding to a cDNA length of 500 to 2000
base pairs was excised and the cDNA was eluted and cloned in
30 the vector pUC13 which had been cleaved with EcoRI and
dephosphorylated. The DNA was then used to transform
competent cells of E. coli MC1016 (lambda). The resulting
colonies were transferred to Pall membranes (Pall
Biodyne), lysed and denatured with 1.5 M NaCl, 0.5 M NaOH
35 and neutralized with 3 M NaOAc, pH 5.5. Screening of
colonies harboring an insert of HIV-3 was performed under

-19-

1 moderately stringent conditions in a buffer containing 5 X
SSC, 5 X Denhardts solution, 0.2 % SDS, 250 mg/ml denatured
5 plasmid containing the SacI - EcoRI fragment of HIV-1
discussed above. Following hybridization, filters were
washed as follows:

1. 1 hour in 2 X SSC, 0.1 % SDS at room temperature.
- 10 2. 30 minutes in 0.1 X SSC, 0.1 % SDS at room
temperature.
3. 20 minutes in 2 X SSC, 0.1 SDS at 42°C.
4. 20 minutes in 0.1 X SSC, 0.1 % SDS at 42°C.

15 Following autoradiography of the filter, several weakly
positive colonies were identified which were then grown for
analysis. It was expected that the positive signal would
either be due to weak homology with the gag or pol regions
of HIV-1, or due to some sequence homology with the R
20 region of the LTR.

C. Sequences contained in HIV-3 cDNA.

25 A clone carrying the largest insert, which was found to be
906 base pairs in length and is referred to
as iso 70-11, was selected for sequence analysis. A number
of subclones of the insert were prepared by digesting the
insert with various restriction enzymes and subcloning the
30 resulting fragments in the pUC 13 vector. Sequence
determinations were performed according to the
dideoxy-method, described by Sanger, (Proc. Natl. Acad. Sci.
USA 74: 5463-5467, 1977), using a kit purchased from
Boehringer which makes use of 17-mer M13 primers. Sequence
35 analysis of cDNA clone iso 70-11 revealed that the insert
corresponded to the 3' end of the viral genome which

-20-

1 possessed a poly (A) chain at the 3' end.

The HIV-3 retrovirus contains a 3' LTR which is composed of a U3 region as well as an R region. Like the 3' LTR region
5 of HIV-1, clone iso 70-11 contains an AATAAA polyadenylation signal located approximately 23 nucleotides from the 3' end of the R region. Analysis of the HIV-3 sequence revealed approximately 70 % homology with the corresponding 3' LTR sequence of HIV-1 and less than 55 % homology with the
10 corresponding sequence of HIV-2.

Conversely, hybridizations using HIV-1 gag - pol sequences as the labeled probe to detect crosshybridization with HIV-3 RNA revealed no detectable hybridization when the
15 hybridization was carried out under stringent conditions. This again indicates that the viruses are only distantly related and that a distinction can be made between HIV-1 and HIV-3 at the nucleic acid level in the region of the genome encompassing the gag and pol genes. This same labeled probe
20 did, however, hybridize to RNA derived from HIV-1 strain SF4.

In addition, the invention relates to a composition comprising at least one antigen, in particular, a protein or
25 glycoprotein of HIV-3 retrovirus. Such a composition can be used in methods for detecting antibodies and in kits for carrying out such methods.

The HIV-3 virus has proven to be a usable as a source of
30 antigen for detecting antibodies in people who have come into contact with HIV-3. As such, the virus may be grown and concentrated by the methods already described and a lysate prepared by treating the virus with a suitable detergent. A preferred detergent for preparing a total

35

-21-

1 viral lysate is Triton X-100, used at a concentration of 0.5
%. Another preferred detergent is Nonidet P-40 (NP-40),
also used at a concentration of 0.5 %.

5 Alternatively, viral protein may be purified from lysates of
the virus. A preferred method for purifying these proteins
is affinity chromatography. For example, the viral antigens
10 may be separated on a preparative polyacrylamide gel and the
individual antigens eluted in purified form. These may
further be used to raise antisera in, for example, rabbits
which are specific for the individual viral proteins. The
IgG fraction derived from immune rabbit serum can be coupled
to a solid phase such as CNBr-activated Sepharose 4B
15 (Pharmacia) and used to selectively remove individual viral
antigens from viral lysates. These proteins may then be
eluted from the affinity support using a low pH buffer and
further purified using standard chromatographic techniques
of which an example is given by Montelaro et al., J. of
20 Virology (1982) 42: 1029-1030.

The invention relates generally to any composition which can
be used for the diagnosis of HIV-3 infection or for tests
which have a prognostic value. These diagnostic procedures
25 involve the detection of antibody in serum or other body
fluid, which are directed against at least one of the
antigens of HIV-3.

Preferred compositions are viral lysates or purified
antigens which contain at least one of the viral core
30 proteins, p12, p16, and p25 or envelope proteins gp41 or
gp120, or pol gene-derived proteins, such as p31. Especially
preferred compositions are those which simultaneously
contain, by way of example, the following proteins,

- 35 - p25 and gp120
 - p25 and gp41

-22-

1

- p25, gp41 and gp120
- p12, p16 and p25
- p25, p31 and gp120

5

It should be understood however, that the above mentioned compositions are only meant to serve as examples and that the invention relates to all lysates or protein preparations containing one or more of the above mentioned proteins or glycoproteins.

10

The invention also relates to any composition in which either HIV-3 viral lysate is used in combination with similarly prepared proteins derived from HIV-1 and/or HIV-2 for the general diagnosis of infection or contact with human immunodeficiency virus without regard to the absolute identity of the virus being detected. For example, such compositions could consist of a mixture of lysates of HIV-1, HIV-2 and HIV-3 or could consist of the following:

15

20

- core proteins of HIV-1, HIV-2 and HIV-3, and in particular the major core protein of each virus type, homologous to the HIV-3 p25 protein.
- envelope glycoproteins of HIV-1, HIV-2 and HIV-3 and in particular the outer envelope glycoproteins of each virus type, homologous to HIV-3 gp120.
- core proteins of HIV-1, HIV-2 and HIV-3 together with the envelope glycoproteins of HIV-1, HIV-2 and HIV-3, in particular the major core protein of each virus type, homologous to the HIV-3 p25 protein, together with the major outer envelope protein of each virus, homologous to HIV-3 gp120.
- a combination of the core proteins and envelope

30

35

-23-

1 proteins of HIV-1, HIV-2 and HIV-3 and in particular
homologous to the HIV-3 proteins p25 and gp120
respectively and a protein derived from the pol gene
5 of HIV-1, HIV-2 and HIV-3, in particular the proteins
of each virus type homologous to the p31 endonuclease
protein of HIV-3.

10 Furthermore, the invention relates to an antigen providing a
single band in polyacrylamide gel electrophoresis, said
antigen comprising, in common with one of the purified
antigens of HIV-3 retrovirus, an epitope that is recognized
by serum of persons carrying anti-HIV-3 antibodies. The
15 amino acid sequences corresponding to these epitopes can
readily be determined by isolating the individual proteins
either by preparative electrophoresis or by affinity
chromatography and determining the amino acid sequence of
either the entire protein or the fragments produced
enzymatically by trypsin or chymotrypsin digestion or
20 chemically by the procedures described in detail below. The
resulting peptide or polypeptides can subsequently be
sequenced by Edman degradation. The invention relates
therefore to any protein, glycoprotein or peptide, either
derived directly from the virus or produced by cloning any
25 cDNA fragments of the virus in bacterial expression vectors,
or viral expression vectors for the expression of inserted
DNA in mammalian or insect cells, and purifying the
expressed protein by the methods described above.
Furthermore, the invention also relates to synthetic
30 peptides, produced either by Merrifield synthesis or Fmoc
chemistry, which may be subsequently purified to homogeneity
and which contain in their sequences epitopes which are
shared by the natural HIV-3 antigens.
Antigens which share epitopes with viral proteins may easily
35 be recognized by their reaction with antibodies present in
the serum of individuals infected with HIV-3, either by

-24-

1 Western blotting, or radioimmunoprecipitation. In the case
of small peptides which are not able to bind to
nitrocellulose, these peptides can be detected by binding to
5 nylon membranes (Pall Biodyne or Amersham) and reacting the
membrane with anti-HIV-3 antiserum. In particular, the
invention relates to epitopes contained in any of the HIV-3
core proteins, p12, p16 and p25 or in a protein which may
contain as part of its polypeptide chain epitopes derived
10 from a combination of the core proteins. Furthermore, the
invention relates to epitopes contained in either of the two
HIV-3 envelope glycoproteins, gp41 and gp120 as well as any
protein which contains, as part of its polypeptide chain,
epitopes derived from a combination of the HIV-3 envelope
15 glycoprotein or a combination of the HIV-3 envelope
glycoproteins and HIV-3 core protein. The invention
additionally relates to polypeptides whose synthesis is
directed by expression vectors constructed by recombinant
DNA methods which incorporate epitopes derived from HIV-3
20 proteins or glycoproteins together with epitopes derived
from the proteins or glycoproteins of either HIV-1 and/or
HIV-2 into a single polypeptide chain. Preparing such a
construction would involve excising the relevant coding
regions from cDNA of HIV-3 as well as HIV-1 and HIV-2, and
25 coupling the DNA in phase so as to form a coding sequence
which, when inserted into an expression vector
possessing the necessary signal sequences, directs the
synthesis of a hybrid protein in which epitopes of the
HIV-3, HIV-1 and HIV-2 are contained.

30 Furthermore, the invention relates to methods for the
detection of antibodies against HIV-3 retrovirus in a
biological fluid, in particular for the diagnosis of a
potential or existing ARC or AIDS caused by HIV-3
35 retrovirus, characterized by contacting body fluid of a
person to be diagnosed with a composition containing one or

-25-

1 more of the proteins or glycoproteins of HIV-3 or with a
lysate of the virus, or with an antigen possessing epitopes
common to HIV-3, and detecting the immunological conjugate
5 formed between the anti-HIV-3 antibodies and the antigen(s)
used.

Preferred methods include, for example, immunofluorescence
assays or immunoenzymatic assays.

10 Immunofluorescence assays typically involve incubating, for
example, serum from the person to be tested with cells
infected with HIV-3 and which have been fixed and
permeabilized with cold acetone. Immune complexes formed
15 are detected using either direct or indirect methods and
involve the use of antibodies which specifically react to
human immunoglobulins. Detection is achieved by using
antibodies to which have been coupled fluorescent labels,
such as fluorsecein or rhodamine.

20 Immunoenzymatic assays may be performed, for example, as
follows:

- 25 - a specific quantity of HIV-3 virus extract or of a
composition referred to according to the invention is
deposited in the wells of a microtitration plate.
- the excess unbound material is removed after a
suitable incubation period by washing.
- 30 - a suitable dilution or dilutions of serum of other
body fluid which is to be tested for the presence of
antibodies directed against one or more of the protein
or glycoprotein antigen of HIV-3 is introduced into
the well.
- the microtitration plate is incubated for a period of
35 time necessary for the binding reaction to occur.
- the plate is washed thoroughly.

-26-

- 1 - the presence of immune complexes is detected using
 antibodies which specifically bind to human
 immunoglobulins, and which have been labeled with an
5 enzyme, preferably but not limited to either
 horseradish peroxidase, alkaline phosphatase, or
 beta-galactosidase, which is capable of converting a
 colorless or nearly colorless substrate into a highly
 colored product. Alternatively, the detection system
10 may employ an enzyme which, in the presence of the
 proper substrate(s), emits light.
- the amount of product formed is detected either
 visually, spectrophotometrically, or luminometrically,
 and is compared to a similarly treated control.

15 Other detection systems which may also be used include those
 based on the use of protein A derived from Staphylococcus
 aureus Cowan strain I, protein G from group C Streptococcus
 sp. (strain 26RP66), or systems which employ the use of the
20 biotin-avidin binding reaction.

 Another method of immunoenzymatic detection of the presence
 of antibodies directed against one or more of the HIV-3
 antigens is the Western blot. The viral antigens are
25 separated electrophoretically and transferred to a
 nitrocellulose membrane or other suitable support. The body
 fluid to be tested is then brought into contact with the
 membrane and the presence of the immune complexes formed is
 detected by the method already described. In a variation on
30 this methods, purified viral antigen is applied in lines or
 spots on a membrane and allowed to bind. The membrane is
 subsequently brought into contact with the body fluid to be
 tested and the immune complexes formed are detected using
 the previously described techniques.

35 The presence of antibodies in body fluid may also be

-27-

1 detected by agglutination. HIV-3 lysates or a HIV-3 lysate,
antigen or purified antigen composition referred to
according to this invention, is used to coat, for example,
5 latex particles which form an uniform suspension. When
mixed with serum containing antibodies to the antigen
present, the latex particles are caused to agglutinate and
the presence of large aggregates can be detected visually.

10 The present invention also relates to labeled extracts of
HIV-3 or compositions as previously described. The labeling
can be of any type, such as enzymatic, chemical, fluorescent
or radioactive.

15 Furthermore, the invention relates to a method for detecting
the presence of HIV-3 antigens in body fluids. This may,
for example, be accomplished in the following manner:

- 20 - the IgG fraction of antiserum, derived either from
humans infected with HIV-3 or from animals injected
with an HIV-3 lysate or composition already described,
is placed in the wells of a microtitration plate.
- after a suitable period to allow adsorption, the
excess unbound material is washed away.
- 25 - a body fluid containing the antigen to be detected is
placed in the well.
- the microtitration plate is allowed to incubate for a
suitable period of time to allow binding to occur.
- the plate is then thoroughly washed with a suitable
30 buffer.
- the presence of bound antigen is detected either
directly or indirectly, for example, by using
immunoglobulins which are similarly specific for the
antigen(s) to be detected and which have been labeled,
preferably with one of the aforementioned enzymes.
- 35 - an appropriate substrate is then added and the extent

-28-

1

of reaction is compared to a control in order to measure the amount of antigen present.

5

Furthermore, the invention relates to a kit for the detection of anti-HIV-3 antibodies in biological fluids, comprising an HIV-3 lysate or a composition as referred to above and a means for detecting the immunological complexes formed.

10

In the case of kits designed to detect specific antibodies by immunoenzymatic methods such a kit would include:

15

- an HIV-3 lysate or composition of one of the types already described, preferably in a purified form, and preferably attached to a solid support such as a microtitration plate.
- a conjugate between an enzyme and an immunoglobulin fraction which is capable of binding to the antibodies to be detected, or a conjugate between an enzyme and bacterial protein A or protein G.
- a control antigen which possesses no epitopes which are shared by any human immunodeficiency virus.
- appropriate buffers for performing the assay.
- an appropriate substrate for the enzyme.

20

25

Kits for the detection of specific antibodies which make use of labeled antigen would include:

30

35

- an appropriately labeled antigen or combination of antigens of the types already described.
- rotein A or anti-human immunoglobulins, preferably coupled to an insoluble support, such as Protein A-Sepharose 4B (Pharmacia) or an equivalent support.
- control antigen, which is not recognized by anti-HIV-3 antisera.

-29-

1

- appropriate buffers for performing the assay.
- if appropriate, substrates for the detection of enzymatically labeled antigen.

5

The invention further relates to kits, developed for the detection of HIV-3 antigens in biological fluids, which comprise :

10

- anti-HIV-3 immunoglobulins, preferably coupled to a solid support such as a microtitration plate.
- anti-HIV-3 immunoglobulins conjugated to an enzyme.
- negative control antigen, which would not be recognized by anti-HIV-3 immunoglobulins.
- positive control antigen which consists of one of the HIV-3 antigens or compositions already described.
- appropriate buffers for conducting the test.
- an appropriate substrate for detection of bound enzyme.

15

20

Furthermore, the invention relates to an immunogenic composition containing an envelope glycoprotein of HIV-3 retrovirus, in particular, gp41 or gp 120, or a part of said glycoprotein, in combination with a pharmaceutically acceptable vehicle suitable for the constitution of vaccines effective against HIV-3. The invention additionally relates to any peptide or polypeptide which contains within its sequence all or part of the protein backbone of the HIV-3 retrovirus, as well as peptides which result from addition, substitution, or deletion of amino acids which do not affect the general immunological properties of said peptides.

25

30

35

The invention further relates to monoclonal antibodies characterized by their ability to specifically recognize epitopes contained in the HIV-3 antigens or compositions as previously defined, and in particular, monoclonal antibodies

-30-

1 raised specifically against said antigens and produced by
traditional techniques. The invention also relates to
monoclonal antibodies of human origin produced by
5 immortalizing B-cells derived from persons infected with
HIV-3, for example, by transforming the B- cells with
Epstein-Barr virus and subcloning the transformants.

10 The invention likewise relates to the production of
polyclonal antisera in animals which recognize one or more
HIV-3 antigens and which is produced by infecting animals
with purified HIV-3 or an HIV-3 antigen or combination of
antigens, and in particular the proteins or glycoproteins of
HIV-3.

15 The antibodies, either polyclonal or monoclonal, can be used
for a wide variety of purposes which include neutralization
of HIV-3 infectivity, the detection of HIV-3 antigens in
biological fluids or in infected cells, and the purification
20 of HIV-3 protein and glycoprotein antigens.

The invention further relates to nucleic acids, optionally
labeled which are derived in part, at least, from RNA of
HIV-3 retrovirus or of variants of this virus.

25 The invention relates likewise to the use of cDNA or parts
of the cDNA or the recombinants containing them, which are
characterized by containing at least a portion of the cDNA
corresponding to the entire genomic RNA of the HIV-3
30 retrovirus. Such cDNAs may be used as probes for the
specific detection of HIV-3 sequences in biological fluids,
tissues and cells. The probes are preferably also labeled,
either radioactively or chemically, alternatively using
enzymatic, fluorescent or chemiluminescent labels which
35 enable the probes to be detected. Preferred probes for the
specific detection of HIV-3 and diagnosis of HIV-3 infection

-31-

1 are probes that contain all or a portion of the cDNA
complementary to the HIV-3 genome. In this context, an
especially advantageous probe can be characterized as one
5 which contains, in particular, the nucleic acid sequence
contained in clone iso 70-11 and which includes the viral
LTR-R sequence which is located at both the 5' and 3' ends
of viral genomic RNA and at both the 5' and 3' ends of
integrated proviral DNA.

10
It is nevertheless understood that the probes which can be
15 used for the diagnosis of HIV-3 infection are in no way
limited to the probes described above, and that the
invention incorporates all sequences which originate from
the HIV-3 genome or its naturally occurring variants and
includes sequences encoding the viral core proteins (gag
20 gene), the two forms of reverse transcriptase and the
endonuclease (pol gene), as well as the two viral envelope
glycoproteins (env gene).

25 The invention also relates to HIV-3 nucleic acid sequences
which have been incorporated into a recombinant nucleic acid
comprising a nucleic acid from a vector, and having said
cDNA or part of said cDNA inserted therein. Such a
construction could be used for replicating the viral cDNA or
its fragments in an organism or cell other than the natural
30 host so as to provide sufficient quantities of the probe to
be used for diagnostic purposes.

A probe generated in such a manner can be employed in a
diagnostic test for specific detection of HIV-3 which
35 incorporates the following essential steps :

-32-

- 1
- labeling of the probe generated as described above by the methods previously described.
 - bringing the probe into contact under stringent
 - 5 hybridization conditions with DNA from infected cells or viral RNA from infected cells or biological fluids, once said DNA or RNA has been, preferably, applied to a membrane and has been rendered accessible to the probe.
 - 10 - washing the membrane with a buffer under circumstances in which stringent conditions are maintained.
 - detection of the labeled probe, preferably by autoradiography in cases in which the probe has been radioactively labeled, or by a suitable
 - 15 immunodetection technique in case the probe has been labeled chemically.

The invention further relates to a process for the production of HIV-3 retrovirus characterized by culturing

20 human T4 lymphocytes or human lymphocytic cell lines of leukemic origin which carry the T4+ phenotype with lymphocytes or cell lines that have previously been infected with an isolate of HIV-3 retrovirus, as well as recovering and purifying the retrovirus from the culture medium.

25 The invention likewise relates to a process for the production of antigens of HIV-3 retrovirus, characterized by lysing the retrovirus, preferably with a detergent, and recovering the lysate containing said antigens.

30 The invention additionally relates to a process for the production of any of the HIV-3 proteins or glycoproteins p12, p16, p25, p31, gp41, gp 120 or reverse transcriptase as previously defined, or a part thereof, characterized by inserting the nucleic acid encoding the proteins or

35 glycoproteins in an expression vector, transforming a host with said vector, culturing the transformed host as well as

-33-

1 recovering and purifying the expressed protein. The process
includes vectors which may or may not direct the synthesis
of fusion proteins and includes but is not limited to
5 bacterial expression vectors, mammalian expression vectors
such as vaccinia virus, and vectors based on baculovirus for
the expression of cloned genes in insect cells.

10 EXAMPLES

MATERIALS AND METHODS

Virus and cell culture

15 a. Virus strains and cell lines.

HUT-78 cells chronically infected with ARV-4 (HIV-1 SF4),
originally isolated by J. Levy, San Francisco, U.S.A. (20)
and
20 uninfected HUT-78 cells were kindly provided by S. Sprecher,
Brussels, Belgium. LAV-2rod originally from L. Montagnier,
Paris, and CEM cells were obtained from J. De Smeyter,
Leuven, Belgium. Isolate 53, an HIV-2 isolate, was isolated
in this laboratory (21).

25 b. Virus isolations.

Virus isolations were performed in a manner similar to that
described by Levy and Shimabukuro (22), with modifications.
30 Lymphocytes from patients as well as from healthy donors
were isolated from heparinized whole blood on Lymphoprep
(Nyegaard and Co., Oslo, Norway) and were cultured in RPMI
1640 containing 20 mM HEPES, 15 percent fetal calf serum
(Gibco), 5 g/ml hydrocortisone (Merck), 75 U/ml IL-2 and 2
35 g/ml polybrene (Aldrich).

-34-

1 Lymphocytes from healthy donors were stimulated with 2 g/ml
phytohemagglutinin (PHA, Wellcome) for 3 days prior to use.
Fresh PHA-stimulated lymphocytes were added to the virus
5 isolation cultures every 3 to 4 days. Cultures were
monitored for cytopathic effect, immunofluorescence, using a
broad specificity, polyclonal reference antiserum (23), and
the presence of antigen in the culture supernatants
(Innotest VCA- HIV, Innogenetics). The broad specificity
10 reference (BSR) anti-serum used was derived from an
HIV-1-infected donor and was shown experimentally to have an
exceptionally high titer ($\geq 1,000,000$ in an enzyme
immunoassay based on recombinant HIV-1 p24 protein) and to
crossreact strongly with the gag and pol gene products of
15 other HIV types, in particular, HIV-2. Reverse
transcriptase was also assayed essentially as described
(24).

In order to establish chronically infected, permanent cell
lines, virus-infected primary lymphocytes were co-cultured
20 with Molt 4 clone 8 cells (25), kindly provided by N.
Yamamoto, Yamaguchi, Japan, and monitored for cell growth.
Virus production was monitored by the reverse transcriptase
assay as well as antigen capturing.

25 Differential antigen capturing.

A test system was developed whereby a distinction can be
made between HIV-1 and other related human immunodeficiency
viruses. The system is based on a comparison of the ability
30 of two different polyclonal IgG preparations, one with a
broad anti-HIV specificity which is due its exceptionally
high titer, particularly against the major core protein, and
one with a lower titer which reacts preferentially with
HIV-1, to capture detergent-treated virus in culture
35 supernatants. Detection of captured antigen is achieved by
using a (broad specificity IgG)/horseradish peroxidase

-35-

1 conjugate.

The test detects primarily but not exclusively the p24 core protein.

5

Monoclonal antibodies to HIV-1

The panel of monoclonal antibodies used has been described (26). The antibodies were prepared against native viral proteins in Triton X-100-disrupted HIV-1 preparations.

10

Protein analysis

a. Electrophoresis.

15

Polyacrylamide gel electrophoresis of viral proteins was performed essentially as described by Maizel (27).

b. Protein blotting.

20

Blotting was performed either in a Bio-Rad transblot cell at 400 mA for 4 hours using the carbonate buffer described by Dunn (28) or using the LKB semi-dry blotting apparatus at 0.8 mA/cm² for 1 hour in 48 mM Tris, 39 mM glycine, 0.0375 % sodium dodecylsulfate (SDS) and 20 % methanol.

25

c. Generation of partial cleavage products.

Viral proteins were analyzed by the technique shown in Figure 1. Advantage was taken of the fact that corresponding proteins from the various isolates have similar molecular weights. Proteins were separated on 12.5 percent SDS- polyacrylamide gels together with a marker lane of ARV-4 proteins which was excised following electrophoresis, blotted and incubated with an anti-HIV antiserum to reveal the positions of the viral proteins.

30

35

-36-

1

The marker blot was in turn used to locate the approximate positions in the Coomassie blue stained portion of the gel of the viral proteins to be cleaved. Horizontal gel slices containing the proteins were excised, transferred to glass tubes and subjected to chemical cleavage.

5

1. Cyanogen bromide cleavage

10

The gel slice was incubated with 10 ml of a freshly prepared 40 mg/ml solution of CNBr (Merck) in 0.3 N HCl for 3 hours at room temperature in a fume hood. Following the incubation, the gel slice was equilibrated with SDS-sample buffer for electrophoresis in the second dimension.

15

2. BNPS-Skatole cleavage.

20

The gel slice was incubated with 10 ml of a freshly prepared saturated solution of 2-(-2'-nitrophenylsulfenyl)-3-methyl-3'-bromoindoline (BNPS-Skatole, Pierce) in 70 percent acetic acid; 30 % H₂O containing 0.1 % phenol, for 3 hours at room temperature, protected from light. Following the incubation, the gel slice was equilibrated by repeated washing in SDS-electrophoresis sample buffer.

25

30

Following cleavage, the individual lanes were excised from the gel slices, rotated 90° and placed on top of a 10 to 20 percent SDS-polyacrylamide gradient gel. On completion of electrophoresis, the gel was blotted onto nitrocellulose (Schleicher and Schuell) and blocked with PBS containing 1 mg/ml casein (Merck). Only cleavage products with molecular weights in excess of 10 kD are able to be visualized since peptides with lower molecular weights do not bind efficiently to nitrocellulose. Blots were incubated with a broad spectrum anti-HIV antiserum followed by goat anti-human IgG: alkaline phosphatase conjugate (Promega).

35

-37-

1 Partial cleavage products were then visualized by reaction
with 5-bromo-4-chloro-3-indolylphosphate and nitro blue
tetrazolium (Sigma).

5 Viral Nucleic Acids

10 a. Hybridization to viral RNA.

Virus from culture supernatants was harvested by pelleting
through cushions of 20 % sucrose by centrifugation at 26,500
rpm for 1,5 hrs. at 4°C and was disrupted in 10 mM Tris, pH
15 7.4, 10 mM NaCl, 10 mM EDTA containing 0,5 % sodium
dodecylsulfate. Aliquots of the disrupted virus were
spotted onto a membrane of Hybond H (Amersham) in amounts
corresponding to 5, 2.5, 1.25 and 0.62 milliliters of
original culture supernatants. The RNA deposited onto the
20 filter was fixed to the membrane by irradiation with
ultraviolet light for 2 hrs. The RNA bound to the filter
was then subjected to hybridization with an HIV-1 cDNA probe
which had been labeled by nick translation with 32p-dCTP.
The hybridization was carried out under stringent conditions
25 in 3 X SSC, 0,5 % milk powder, 1 % SDS, 10 % dextran sulfate
and 50 % formamide at 42°C for 18 hrs. Following
hybridization, the filter was washed twice under stringent
conditions in 0.1 X SSC and 0.1 % SDS for 30 minutes.
Hybridization was detected by autoradiography at -70°C with
30 enhancing screens. Hybridizations were similarly performed
using a probe derived from the env region of HIV-2.

Hybridizations were also performed under nonstringent
conditions in 5 X SSC, 25 % formamide, 5 X Denhardt's
35 solution, 10 % dextran sulfate, and 100 g/ml denatured
salmon sperm DNA at 37°C overnight. The filter was

-38-

1 subsequently washed 4 times for 15 minutes in 5 X SSC, 0.1 % SDS at room temperature and autoradiographed.

5 b. Preparation of ANT 70 cDNA

10 Virus was pelleted from 1 liter of culture supernatant using polyethylene glycol 6000, redissolved in PBS and pelleted through a 20 % sucrose cushion. The resulting pellet of virus was disrupted in 6 M guanidinium chloride in 20 mM sodium phosphate buffer, pH 6.5, containing 20 mM dithiotreitol and 0.5 % NP-40. Solid CsCl was added to a concentration of 2 molar. The solution containing disrupted virus was layered onto a cushion of 5.7 M CsCl containing
15 0.1 M EDTA and the viral RNA was pelleted by centrifugation at 25,000 in a Beckman SW 28 rotor at 15°C for 20 hrs. Following centrifugation, the RNA was redissolved, extracted with phenol and precipitated with ethanol and 2 M LiCl.

20 One-fifth of the viral RNA prepared was used to direct the first step in the synthesis of cDNA using a kit supplied by Amersham. cDNA synthesis was primed using oligo (dT). the synthesis was carried out using the reverse transcriptase supplied with the kit. Second strand synthesis was
25 performed using E. coli DNA polymerase I in the presence of RNase H to digest away the RNA strand of the RNA/DNA hybrid. The synthesis of the second strand was performed in the presence of 32P-dCTP to label the cDNA. The resulting cDNA was treated with T4 DNA polymerase to create blunt ends,
30 the cDNA was methylated to protect possible internal EcoRI cleavage sites, and was then coupled to EcoRI linkers (Amersham). The EcoRI sites in the linkers were then cleaved and the cDNA was sized on a 1.2 % agarose gel. The region of the gel corresponding to a cDNA length of 500 to
35 2000 base pairs was excised, and the cDNA was eluted and cloned in the vector pUC13 which had previously been cleaved

-39-

1 with EcoRI and dephosphorylated. After ligation, the DNA
was used to transform competent cells of E. coli MC1016
(lambda). The resulting colonies were transferred to Pall
5 membrane filters (Pall Biodyne), lysed and denatured with
1.5 M NaCl, 0.5 M NaOH and neutralized with 3 M NaOAc, pH
5.5. Screening of colonies harboring an insert of HIV-3 was
carried out by hybridization under moderately stringent
conditions in 5 X SSC, 5 X Denhardt's solution, 0.2 % SDS,
10 250 mg/ml denatured salmon sperm DNA overnight at 65°C.
Hybridization was performed using the HIV-1 SacI-EcoRI
fragment. Following hybridization, the filters were washed
as follows:

- 15 1. 1 h. in 2 X SSC, 0.1 % SDS at room temperature.
2. 30 minutes in 0.1 X SSC, 0.1 % SDS at room temperature.
3. 20 minutes in 2 X SSC, 0.1 % SDS at 42°C.
4. 20 minutes in 0.1 X SSC, 0.1 % SDS at 42°C.

20 After washing, the filters were autoradiographed at -70°C
using intensifying screens.

Hybridizations were also performed under the nonstringent
conditions used for nonstringent hybridization of the HIV-1
25 and HIV-2 probe.

c. Analysis of cDNA clones.

30 Colonies giving a positive hybridization signal were grown
for analysis. Plasmids were isolated, cleaved with EcoRI
and subjected to agarose gel electrophoresis to confirm the
presence of an insert and to determine its size. Of 96
colonies analyzed 17 were found to contain inserts. Five
were taken for further analysis and ranged in size from
35 approximately 800 to 1600 base pairs in length.

-40-

1

d. Sequence determinations.

5

Nucleotide sequence determinations were performed according to the dideoxynucleotide method of Sanger (Proc. Natl. Acad. Sci. USA 74: 5463-6467, 1977), using a kit supplied by Boehringer. Sequencing was carried out using 17-mer M13 primers.

10

e. Hybridizations of ANT 70 cDNA to HIV-1 and HIV-2 viral RNA.

15

The ANT 70 cDNA clone containing the largest insert (iso 70-11) was used for hybridization to the filter onto which viral RNAs had been deposited.

20

Hybridization was performed under stringent conditions in 3 X SSC, 0.5 % milk powder, 1 % SDS, 10 % dextran sulfate, and 50 % formamide at 42°C for 18 hrs. Following hybridization, the filter was washed with 0.1 X SSC, 0.1 % SDS at 65°C (2-30 minute washes) after which the filter was autoradiographed at -70°C with an intensifying screen.

25

30

35

-41-

1 Results

Virus isolation

5 As part of a continuing study on heterosexual transmission of HIV, a virus isolation was performed from blood from a Camerounian woman and her partner. As before, the two isolated strains will be named HIV-3 (ANT 70) (woman) and HIV-3 (ANT 70 NA) (man), respectively. For convenience, the shorter terms ANT 70 and ANT 70 NA will also be used. The
10 woman is the partner of an HIV- seropositive man with generalized lymphadenopathy. Serum from the woman was moderately positive (ratio O.D./cut-off of 4.5) in the enzyme-linked immunosorbent assay (EIA, Organon Teknika) and had a low titer (1/40) in the immunofluorescent antibody
15 assay for HIV-1 but gave ambiguous results in the HIV-1 Western blot assay with clear bands at p33, P53/55 and p64 but very weak bands at p24, gp41 and gp120. The woman had elevated serum IgG and IgM levels and a CD4/CD8 ratio of 0.46. Virus was isolated by co- cultivation of the woman's
20 lymphocytes with PHA-stimulated lymphocytes from healthy uninfected donors. After 52 days in culture, virus was detected in the culture as judged by the presence of syncytia and on the basis of positive immunofluorescence observed when a laboratory reference anti-HIV antiserum was
25 incubated with acetone-fixed cells from the culture. The presence of reverse transcriptase was also detected in the

30

35

-42-

1 culture supernatant (10^4 cpm/ml, 27 X background).
Cell-free culture supernatant was used to passage the virus
on fresh lymphocytes. After 15 days, CPE was again observed
5 and reverse transcriptase detected in the supernatant. A
comparison of detergent-treated culture supernatant from
this isolate (ANT 70) with other isolates by differential
antigen capturing revealed, however, that this isolate was
not HIV-1.

10 These results are illustrated in Figure 2. It is evident by
the lower O.D. values that the isolate (ANT 70) is, in
contrast to the other isolates, poorly recognized by the
HIV-1 specific IgG but, like the other isolates, was
readily captured by the broad specificity IgG (panel F).
15 The other isolates, which were subsequently all shown to be
HIV-1 strains using an HIV-1 specific MAb (CLB MAb 14), all
gave higher O.D. values on the plates coated with specific
IgG than on plates coated with the broad specificity
reference IgG.

20 An attempt was made to transfer the virus to a permanent
cell line by co-cultivating isolate (ANT 70)-infected
primary lymphocytes with Molt-4 clone 8 cells. In the
initial phase of the infection, extensive cytopathic effect
25 was observed with syncytium formation and cell death.
Within several weeks, cell growth was detected. The cells
gave a positive immunofluorescence when tested using a broad
spectrum anti-HIV antiserum and the presence of antigen and
reverse transcriptase was easily detectable in the culture
30 supernatant.

Virus was similarly isolated from the partner of the woman
from whom isolate (ANT 70) was isolated (strain ANT 70 NA).
The man was suffering from lymphadenopathy and was
35 classified as class 3 according to the CDC classification
system. The man also had elevated serum IgM and IgG levels

-43-

1 and a CD4/CD8 ratio of 0.4. Virus was detected in the
supernatant of the culture on day 18. Detergent-treated
supernatant containing this virus was also analyzed by
5 differential antigen capturing and found to react in a
manner similar to isolate (ANT 70) (Figure 3). The binding
of antigen derived from this isolate was again less with
HIV-1 specific IgG than with the broad specificity IgG.

10 Serum from the person from whom the isolate (ANT 70 NA) was
derived was incubated with HIV-1 and HIV-2 Western blot
strips (Biotech). Additional strips were also incubated
with serum from a donor infected with HIV-1 as well as serum
from the person from whom HIV-2 (isolate 53) was isolated.
15 These results are shown in Figure 4. Serum from the person
infected with ANT 70 NA crossreacted to a significant extent
with virtually all HIV-1 proteins, including the envelope
proteins. In contrast, serum from the HIV-2-infected
individual crossreacted only with the gag p24 protein, p34
20 endonuclease and p68 reverse transcriptase. The
anti-HIV-1 serum recognized only the p26 gag protein of
HIV-2, while serum from the carrier of ANT 70 NA recognizes
this protein and the HIV-2 reverse transcriptase.

25 Characterization of viral proteins.

Virus in the culture supernatant was precipitated using
polyethylene glycol 6000 (Merck) and the resulting material
30 was redissolved and pelleted through a 15 percent sucrose
cushion. The pelleted virus was dissociated in SDS-sample
buffer and analyzed by polyacrylamide gel electrophoresis
followed by protein blotting. The blot, shown in figure 5,
was incubated with a broad specificity anti-HIV serum to
35 reveal the viral proteins.

-44-

1 In addition to reacting with all of the HIV-1 viral
proteins, the BSR antiserum also crossreacts with the gag
and pol gene products of HIV-2. This antiserum clearly
5 recognizes the gag and pol gene products of ANT 70 as well.
It is evident that the molecular weights of the ANT 70 gene
products differ from those of either HIV-1 or HIV-2. The
molecular weights of the various viral proteins are
summarized in table 1. The variability in the HIV-1 p17/p18
10 protein is due to a 6 amino acid insertion which is present
in some strains between positions 120 and 121 in the HIV-1
HXB2 sequence. A comparison of the proteins from ANT 70 and
ANT 70 NA are shown in figure 6. The molecular weights of
all of the proteins of ANT 70 NA are identical to those of
15 ANT 70.

In order to investigate further the antigenic relationship
between HIV-1, HIV-2 and ANT 70, a series of African and
European anti- HIV-1 sera were diluted 1:1000 and used to
20 coat microwell plates for antigen capturing.
Detergent-treated culture supernatant containing HIV-1, ANT
70, HIV-2 (LAV-2rod) and HIV-2 (isolate 53) were diluted and
the ability of each antiserum to capture the four different
isolates was analyzed. Representative results are shown in
25 Figure 7. It can be seen from this experiment that the
ability of the various sera to capture HIV-1 is in no way
related to their ability to capture either HIV-2 or ANT 70.
In contrast, the ability of these sera to capture LAV-2rod,
the prototype HIV-2 strain, is strongly correlated with the
30 ability of these sera to capture isolate 53, which is also
an HIV-2 strain but an independent isolate. These data
indicate that ANT 70 is neither HIV-1 nor HIV-2.

In a series of related antigen capturing experiments, four
African anti-HIV-1 sera were chosen in order to access their
35 ability to bind HIV-1, ANT 70, HIV-2 (LAV-2rod) and HIV-2
(isolate 53) when the IgGs were coated at different

-45-

1 dilutions. Culture supernatants were diluted so as to give
approximately the same optical density when captured on
plates coated with the IgG used in panel B of Figure 8.
5 Dilutions of the four sera were coated and virus- containing
supernatant was added. The assumption was made that similar
viruses should give rise to similar titration curves.
Indeed, in Figure 5, LAV-2rod and isolate 53 both react
similarly with the coated IgGs. On the other hand, ANT 70
10 gave more intense signals at higher IgG dilutions than did
either of the HIV-2 isolates and the shapes of the curves
obtained with ANT 70 resemble more closely the curves
obtained for HIV-1, except that the optical densities are
consistently lower.

15
Cross reactivity of mouse monoclonal antibodies directed
against HIV-1 p24 core protein.

20 A panel of mouse monoclonal antibodies (MAbs) prepared
against the HIV-1 p24 core protein was tested for their
ability to crossreact with ANT 70 and HIV-2 isolates. In
principle, any panel of anti- HIV-1 p24 monoclonal
25 antibodies can be used, as long as the series includes
monoclonal antibodies which react with different epitopes on
the HIV-1 p24 molecule. Ascites fluid containing the
antibodies was diluted and used to coat microwell plates.
Detergent-treated, virus-containing supernatants were then
30 added to the coated wells. Bound antigen was detected using
BSR-HIV IgGs conjugated to horseradish peroxidase. The
results obtained are shown in Figure 9.
In control wells coated with polyclonal broad spectrum IgGs,
all virus-containing supernatants gave optical densities
35 which exceeded the limits of the microwell plate reader.
However, when tested in wells coated with the various

-46-

1

monoclonal antibodies, quite a different pattern emerged. Previous studies indicated that all of the MABs tested react against different epitopes on the p24 molecule with the exception of MABs CLB 59 and CLB 21 which have been shown to recognize the same epitope. Both of these two MABs react strongly with HIV-1 as expected and also give a measurable signal with ANT 70 but fail to react with either of the HIV-2 strains. Two other MABs, CLB 64 and CLB 14, bound HIV-1 well and showed a weak affinity for ANT 70 as well as the two HIV-2 isolates. In particular, MAB CLB 14 has been shown to recognize all HIV-1 isolates well (> 150 tested). This MAB must therefore bind to a very highly conserved epitope, remnants of which can also be detected in other human immunodeficiency viruses. The other MABs to p24 (CLB 16, 47 and 19.7) and two others which were raised against the HIV-1 p18 protein (CLB 13.4 and CLB 13.6), failed to recognize either ANT 70 or the two HIV-2 isolates but did capture the corresponding HIV-1 antigens.

20

Reaction of human anti-HIV antisera to viral proteins.

Protein blots of viral proteins from HIV-1 (ARV 4), ANT 70 and HIV-2 (LAV2rod) were prepared after electrophoresis of detergent-solubilized extracts and incubated with various human sera (Figure 10). Panel A shows the reaction of the broad specificity laboratory reference serum with the three virus isolates. In panel B, an anti-HIV-1 antiserum was incubated with the blot and recognizes preferentially HIV-1 proteins. Serum from the woman from whom ANT 70 was isolated (panel C) and her partner from whom ANT 70 NA was isolated (panel D) were tested for their ability to recognize other viral isolates. Both of these sera preferentially recognize ANT 70 including the gp120 envelope protein of this virus. Serum from the partner has a higher titer than serum from the woman from whom ANT 70 was

-47-

1 isolated and recognizes the gp41 of HIV-1. Both of these
sera have a higher affinity for ANT 70 than for HIV-1 or the
HIV-2 isolates. In contrast, serum from the person from
5 whom HIV-2 isolate 53 was isolated binds preferentially to
HIV-2 proteins and recognizes the HIV-2 gp120 envelope
protein of this virus as well as the gp41 transmembrane
protein (panel E). It does not react with glycoproteins of
HIV-1 or ANT 70. These results further indicate that ANT 70
10 is different from either HIV-1 or HIV-2.
Enzyme immunosorbent assays using coated viral proteins
titrations of anti-HIV-1, anti-ANT 70 and anti-HIV-2 sera
were performed in microwell plates coated with HIV-1
(ARV-4), ANT 70 and HIV-2 (isolate 53) viral lysates.
15 Two-fold dilutions of each sera, beginning at an initial
dilution of 1:100, were tested for their ability to bind to
the coated antigen. Bound antibody was detected using a
horseradish peroxidase-labeled goat anti-human IgG
conjugate. These results are shown in Figure 11. The anti-
HIV-1 serum recognized preferentially the HIV-1 proteins but
20 shows a significant amount of crossreaction with ANT 70
proteins. The HIV-2 proteins were barely detected. In
contrast, anti-ANT 70 serum preferentially recognized ANT 70
proteins, showed crossreactivity toward HIV-1 proteins, and
reacted better with the HIV-2 coated wells than did the
25 anti-HIV-1 serum as evidenced by the higher optical density
values obtained. The anti-HIV-2 serum had a very low titer
but nevertheless reacted best with HIV-2 proteins. No
detectable signal was observed on HIV-1 or ANT 70 coated
wells. The inability to detect crossreaction in this
30 instance is undoubtedly related to the low anti-HIV titer of
this serum.

-48-

1

Analysis of partial chemical cleavage products of viral proteins.

5

10

15

20

The two reagents used for chemical cleavage, cyanogen bromide and BNPS-skatole, were chosen because of their high specificities for methionine (29) and tryptophan (30), respectively. These two amino acids are also rather hydrophobic and are therefore also less likely to found located in epitopes on the outer surfaces of protein molecules (31). Examination of published amino sequences of the gag and pol gene products of HIV-1 (32-36), HIV-2 (19), SIVagm (10), SIVmac (9), equine infectious anemia virus (EIAV, 37) and Visna (38) reveals that while there is little amino acid homology between some of these diverse isolates, many of the positions of the methionine residues in these proteins and, to an even greater extent, the tryptophan residues, are strikingly conserved (figure 12). Furthermore, intraspecies variation in these residues is minimal or absent, at least in the case of HIV-1 (36) and probably holds true for all of the human and simian immunodeficiency retroviruses.

25

30

35

The partial digestion patterns of the gag and pol gene products of HIV-1, ANT 70, HIV-2 (LAVrod) and HIV-2 (isolate 53) are shown in Figure 13.

Inspection of the CNBr cleavage patterns of the p24 protein from the four isolates reveals that the patterns generated for HIV-2 (LAV-2rod) and HIV-2 (isolate 53) are identical. Different patterns, however, are observed for HIV-1 and for ANT 70. Thus, significant differences exist in the locations of the methionine residues in the major core protein of HIV-1, ANT 70 and HIV-2. In the case of the p17 core protein, differences are observed between the two HIV-2 isolates. Inspection of the published sequence for HIV-2rod

-49-

1 indicates that there is a methionine located 18 amino acids
from the carboxyl terminus of this protein. We conclude
that this methionine must be absent in the corresponding
5 protein from isolate 53. From the cleavage pattern it is
also possible to deduce the presence of a methionine near
(10-15 amino acids) one of the termini of the p16 from ANT
70. CNBr cleavage of the retroviral reverse transcriptase
reveals that again, the proteins from the two HIV-2
10 isolates are identical, while different patterns are
observed for both HIV-1 and ANT 70 proteins. In the case of
the p31 endonuclease derived from the 3'-portion of the pol
gene, similarities can be deduced between all of the
isolates although some minor differences are apparent.

15 BNPS-skatole cleavage of the p24 proteins from the four
isolates results in strikingly similar patterns. It is
evident from Figure 8 that this is to be expected since the
tryptophan positions in this protein are very highly
20 conserved, particularly for the retroviruses of human and
simian origin. We conclude that the tryptophan positions in
the ANT 70 p25 protein also conform to this pattern. .
Inspection of the patterns reveals, however, that minor
differences can be observed, not in the overall appearance
25 of the pattern but rather in the apparent molecular weights
of the species generalized by cleavage. In particular,
differences are detected in the apparent molecular weights
of the central spots in each pattern. As expected, the
patterns for HIV-2 (LAV-2rod) and HIV-2 (isolate 53) are
30 identical. The central spot in the pattern for ANT 70 has
however, a larger apparent molecular weight while the
central spot for HIV-1 (ARV-4) has a lower molecular weight.
In regard to the p16, the positions of the tryptophans in
the ANT 70 protein appear to resemble more closely the
35 positions of the tryptophans found in the HIV-2 protein.
The HIV-1 p17 has a tryptophan located 16 amino acids from

-50-

1

5

the amino terminus of the protein and gives rise to an additional spot not seen in the ANT 70 and HIV-2 patterns following BNPS-skatole cleavage. The tryptophan corresponding to the one at position 36 in the HIV-1 p17 sequence is conserved in all isolates.

10

15

The patterns generated by cleavage of the reverse transcriptase from the four isolates are complex but is is once again apparent that the two HIV-2 isolates are identical. Patterns are obtained for ANT 70 which corresponds neither to the pattern obtained for HIV-1 nor to the HIV-2 pattern. Differences in apparent molecular weights of the cleavage products of the p31 endonuclease are also observed but the patterns generated from the corresponding proteins from HIV-1, ANT 70, and HIV-2 also show common features which suggests a conserved structure.

Results

20

Viral Nucleic Acids

a. Hybridization of HIV-1 and HIV-2 cDNA to viral RNAs.

25

30

35

Nucleicacids crosshybridization between HIV-1 and RNA from the viruses HIV-2 and ANT 70 was evaluated by performing the hybridization with the SacI-BglII HIV-1 restriction fragment which had been inserted into the vector pUC13. This fragment contains a portion of the 5' LTR, including the R region, the entire gag gene and most of the pol gene of HIV-1. Under stringent hybridization conditions, hybridization was only observed between this probe and the RNA derived from HIV-1 (SF4). No hybridization was observed between the probe and either HIV-2 or ANT 70 (figure 14). This indicates that the gag and pol regions of HIV-2 and ANT 70 are significantly different from the corresponding region

-51-

1

of HIV-1.

5

The HIV-2 probe used contains a sequence of approximately 1000 base pairs derived from the env gene of HIV-2. This probe hybridized only to HIV-2 RNA under stringent hybridization conditions and no hybridization was observed with either HIV-1 or HIV-3.

10

b. Homology between ANT 70 cDNA and sequences of HIV-1 and HIV-2.

15

The cDNA clone iso 70-11 was used as a probe to assess the degree of nucleic acid homology between the various virus isolates. The filter onto which aliquots of HIV-1, HIV-2 and ANT 70 had been deposited was subjected to hybridization under stringent conditions. The results are also shown in Figure 14. The experiment demonstrates that under stringent hybridization conditions, no crosshybridization can be detected between any of the virus isolates. The ANT 70 derived probe hybridizes only to ANT 70.

20

25 c. Sequence analysis of clone iso 70-11.

Subclones of the insert were made in pUC13 and sequenced using the dideoxynucleotide method.

30 The presence of a poly A tail confirmed that the iso 70-11 insert is derived from the 3' end of the viral RNA.

Adjacent to the poly A tail is the sequence corresponding to the R region of the viral 3' LTR. Sequence contained in the ANT 70 cDNA and the viral sequences to which they correspond

35 are shown below:

-52-

1

1. HIV-3 LTR

	10	20	30	40	50	60
5	CCCATGGATT	TGAAGATACA	CATAAAGAAA	TACTGATGTG	GAAGTTTGAT	AGATCTCTAG
	70	80	90	100	110	120
	GCAACACCCA	TGTTGCTATG	ATAACTCACC	CAGAGCTCTT	CCAGAAGGAC	TAAAAACTGC
	130	140	150	160	170	180
	TGACCTGAAG	ATTGCTGACA	CTGTGGAACT	TTCCAGCAAA	GACTGCTGAC	ACTGCGGGGA
10	190	200	210	220	230	240
	CTTCCAGTIG	GGAGGGACAG	GGGGGGGTTC	GGGGAGTGGC	TAACCTCAG	AAGCTGCATA
	250	260	270	280	290	300
	TAAGCAGCOG	CTTCTGCTT	GTACCGGGTC	TOGGTTAGAG	GACCAGGTCT	GAGCCCGGGA
		U3	←→	R		
15	310	320	330	340	350	360
	GCTCCCTGGC	CTCTAGCTGA	ACCGCTOGT	TAACGCTCAA	<u>TAAAGCTTGC</u>	CTTGAGTGAG

Polyadenylation signal

A - POLY A

←

20

25

30

35

-53-

1

2. Sequences from HIV-3 3'ORF

5 10 20 30 40 50 60
AACATGGGAAACGCATTGAGAAAAGGTAAATTTGAGGGATGGGCACCGTAAGAGAAAGA
AsnMetGlyAsnAlaLeuArgLysGlyLysPheGluGlyTrpAlaAlaValArgGluArg
70 80 90 100 110 120
ATGAGAAGAACTAGAACTTTCCCTGAGTCTGAAOCCATGOGCACCTGGAGTAGGACAGATC
10 MetArgArgThrArgThrPheProGluSerGluProCysAlaProGlyValGlyGlnIle
130 140 150 160 170 180
TCCAGGGAATTAGCAGCTAGAGGAGGGATACCAAGTTCCCATACTCCTCAAAACAATGCA
SerArgGluLeuAlaAlaArgGlyGlyIleProSerSerHisThrProGlnAsnAsnAla
190 200 210 220 230 240
15 GCCCTTGCATTCTAGAAAGTCACCAAGAGGAAGAAGTAGGTTTTCCAGTAGCACCTCAA
AlaLeuAlaPheLeuGluSerHisGlnGluGluGluValGlyPheProValAlaProGln
250 260 270 280 290 300
GTGCTCTAAGGCCAATGACCTATAAAGGAGCATTTGACCTCAGCTTCTTTTTAAAGAA
ValProLeuArgProMetThrTyrLysGlyAlaPheAspLeuSerPhePheLeuLysGlu
310 320 330 340 350 360
20 AAGGGAGGACTGGAAGGGTTAATTTACTCCCATAAAGAGCAGAAATCCTGGATCTTTGG
LysGlyGlyLeuGluGlyLeuIleTyrSerHisLysArgAlaGluIleLeuAspLeuTrp

GTGTATAA
25 ValTyr

30

35

-54-

1 Discussion

We have isolated a novel human immunodeficiency-associated retrovirus from a Camerounian woman (ANT 70) and her partner
5 (ANT 70 NA). At the time the original virus isolation was performed, the woman was only slightly seropositive, gave ambiguous results in the western blot test and was clinically asymptomatic. Since that time, the woman has begun to develop some of the symptoms of AIDS-related
10 complex (ARC). In contrast, her partner, from whom we were also able to isolate a virus with the same characteristics as the original isolate, was suffering from lymphadenopathy and has since developed other symptoms characteristic of AIDS. This novel isolate may therefore be considered to be
15 a human immunodeficiency virus. The fact that this same virus could be isolated from sexual partners also suggests a mode of transmission which is similar to that of human retroviruses.

20 The virus was first recognized as being different from HIV-1 on the basis of its altered ability to be captured in a differential antigen capturing assay. This has proven to be a highly reliable test which is able to distinguish between HIV-1 and non-HIV-1 strains. That this isolate is not HIV-1
25 is borne out at the protein level by 1.) the differing molecular weights of the viral proteins, 2.) a different pattern of crossreactivity with anti- HIV-1 antiserum than HIV-1, 3.) a drastically reduced ability to be recognized by mouse monoclonal antibodies raised against HIV-1 p24 and p17
30 core proteins, 4.) preferential recognition of ANT 70 proteins over HIV-1 proteins by antisera from the virus carrier, and 5.) patterns of partial cleavage of four of the most highly conserved viral proteins which do not match the patterns obtained when HIV-1 proteins are subjected to
35 the same treatment. Nevertheless, sera from the two individuals infected with this virus recognize the HIV-1 gp41 envelope protein. By the same criteria listed above, it is also clear that ANT 70 is not HIV-2. Indeed, the

-55-

1 antigenic differences between ANT 70 and HIV-1 are
smaller than those between HIV-2 and HIV-1. This is
particularly evident from the results presented in Figures 8
and 10.

5

Additional compelling evidence that ANT 70 is a unique
virus different from HIV-1 and HIV-2 comes from the partial
peptide maps. We have shown that there are significant
differences in the most highly conserved viral proteins.

10 The two HIV-2 isolates which were used for comparison gave
essentially identical cleavage patterns except in the case
of CNBr cleavage of the p17 core protein. It should be
noted, however, that the p17 core protein exhibits more
variability than the p24 protein, at least in HIV-1 strains
15 (34). Whether or not this also holds true for HIV-2 awaits
sequence determination on more strains than have been
analyzed to date.

In light of the fact that ANT 70 is antigenically more
closely related to HIV-1 than is HIV-2, as evidenced by a
20 higher degree of crossreactivity which extends even to the
gp41 envelope protein, it was essential to establish that
ANT 70 was more than simply a genetic variant of HIV-1.
This was possible by investigating the locations of some of
the most highly conserved amino acids in a number of viral
25 proteins which are least subject to genetic variation. That
major differences were noted in the cleavage patterns
indicates that HIV-1, HIV-2 and ANT 70 are three genetically
distinct viruses. On the other hand, the same series of
experiments also revealed similarities between the viruses
30 which may indicate that all three arose from a common
progenitor.

The hybridization data also support the notion that ANT 70
is fundamentally different from either HIV-1 and HIV-2. As
35 long as the conditions under which the hybridization is
performed are stringent, a distinction can easily be made
between the three virus types.

-56-

1 Analysis of the cDNA sequences revealed that the insert is
derived from the 3' end of the viral genome. An analysis of
the homology between these sequences and the sequence of
HIV-1 and HIV-2 reveal that ANT 70 is somewhat more closely
5 related to HIV-1, particularly in the LTR sequences (approx.
70 % homology). The differences are nevertheless of such
magnitude as to rule out the possibility that ANT 70 is
simply a genetic variant of HIV-1. The ANT 70 3'LTR also
contains the signal sequences which are typical of
10 retroviral LTRs.

The existence of a third type of human immunodeficiency
virus has immediate epidemiological implications and
consequences for blood bank testing. As has been shown,
15 antibodies from people infected with this virus react
preferentially with this virus, although these antibodies
also crossreact with HIV-1 proteins. While it was possible
to detect a positive reaction of ANT 70 NA serum in enzyme
immunoassays, immunofluorescence assays and Western Blot
20 assays based on HIV-1 proteins, the fact that the positive
signal was due to a crossreaction inevitably implies that
the sensitivity of such tests will be less for antibodies
produced in response to this virus. This was amply
demonstrated by the enzyme immunoassay results (Figure 11).
25 Furthermore, one criterion for seropositivity in the Western
blot assay is the presence of detectable antibodies to both
a gag and/or pol protein and one of the envelope proteins.
Since in the case of the two individuals infected with ANT
70 and ANT 70 NA, respectively, crossreaction was observed
30 to both HIV-1 p24 and the envelope proteins, the conclusion
which is invariably drawn is that these individuals are
infected with HIV-1 but for some reason fail to develop high
titers against HIV-1. It is possible therefore, that this
virus is more widespread than is currently realized. From
35 an epidemiological standpoint, it is essential to develop

-57-

1

specific diagnostic tests for this virus in order to
evaluate the limits of the geographical area in Africa in
which the virus can be found, and to evaluate the extent to
which this virus has been disseminated.

5

10

15

20

25

30

35

-58-

1

REFERENCES

5

1. Dalgleish, A.G., Beverly, P.C.L., Clapham, P.R., Crawford, D.H., Greaves, M.F. and Weiss, R.A. (1984). The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. Nature 312: 763-766.

10

2. Maddon, P.J., Dalgleish, A.G., McDougal, J.S., Clapham, P.R., Weiss, T.A. and Axel, R. (1986). The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. Cell 47: 333-348.

15

3. Wong-Staal, F. and Gallo, R.C. (1985). Human T-lymphotropic retroviruses. Nature 317: 395-403.

20

4. Clavel, F., Guetard, D., Brun-Vezinet, F., Chamaret, S., Rey, M.A., Santos-Ferriera, M.D., Laurent, A.G., Dauguet, C., Katlama, C., Rouzioux, C., Klatzmann, D., Champalimaud, J.L. and Montagnier, L. (1986). Isolation of a new human retrovirus from West-African patients with AIDS. Science 233: 343-346.

25

30

5. Albert, J., Bredberg, U., Chiddi, F., Bottinger, B., Fenyo, E.M., Norrby, E. and Biberfeld, G. (1987). New pathogenic human retrovirus of West-African origin (SBL 6669) and its relationship to HTLV-IV, LAV-II and HTLV-IIIB. AIDS Res.

35

-59-

1 6. Kanki, P.J., Kurth, R., Becker, W., Dreesman, G.,
McLane, M.F. and Essex, M.
Antibodies to simian T-lymphotropic virus type III in
5 African green monkeys and recognition of STLV-III viral
protein by AIDS and related sera.
Lancet 1985, i: 1330-1332.

10 7. Kanki, P.J., Alroy, J. and Essex, M. (1985).
Isolation of T-lymphotropic retrovirus related to HTLV-
III/LAV from wild-caught African green monkeys.
Science 230: 951-954.

15 8. Daniel, M.D., Letvin, N.L., King, N.W., Kannagi, M.,
Sehgal, P.K., Hunt, R.D., Kanki, P.J., Essex, M. and
Desrosiers, R.C. (1985).
Isolation of T-cell tropic HTLV-III-like retrovirus from
macaques.
Science 228: 1201-1204.

20 9. Chakrabarti, L., Guyader, M., Alizon, M., Daniel, M.D.,
Desrosiers, R.C., Tiollais, P., and Sonigo, P. (1987).
Sequence of simian immunodeficiency virus from macaque and
its relationship to other human and simian retroviruses.
25 Nature 328: 543-547.

30 10. Franchini, G., Gurgio, C., Guo, H.-G., Gallo, R.C.,
Collati, E., Fargnoli, K.A., Hall, L.F., Wong-Staal, F. and
Reitz, M.S. Jr. (1987).
Sequence of simian immunodeficiency virus and its
relationship to the human immunodeficiency viruses.
Nature 328: 539-543.

35

-60-

- 1
11. Benn, S., Rutledge, R., Folks, T., Gold, J., Baker, L.,
McCormick, J., Feorino, P., Piot, P., Quinn, T., and Martin,
M. (1985)
5 Genomic heterogeneity of AIDS retroviral isolates from North
America and Zaire.
Science 230: 949-951.
- 10 12. Hahn, B.H., Shaw, G.M., Taylor, M.E., Redfield, RR.,
Markham, P.D., Salahuddin, S.Z., Wong-Staal, F., Gallo,
R.C., Parks, E.S. and Parks, W.P. (1986).
Genetic variation in HTLV-III/LAV over time in patients with
AIDS or at risk for AIDS.
Science 232: 1548-1553.
- 15 13. Magasiny, S., Spire, B., Barré-Sinoussi, F. and
Chermann, J.- C. (1986).
Genomic variability of selected LAV-related AIDS
retroviruses.
20 AIDS Res. 2: 19-30.
- 25 14. Alizon, M., Wain-Hobson, S., Motagnier, L. and Sonigo,
P. (1985).
Genetic variability of the AIDS virus: nucleotide sequence
analysis of two isolates from African patients.
Cell 46: 63-74.
- 30 15. Starcich, B.R., Hahn, B.H., Shaw, G.M., McNeely, P.D.,
Modrow, S., Wolf, H., Parks, E.S., Parks, W.P., Josephs,
S.F., Gallo, R.C., Wong-Staal, F. (1986).
Identification and characterization of conserved and
variable regions in the envelope gene of HTLV-III/LAV, the
retrovirus of AIDS. Cell 45: 637-648.
- 35

-61-

1 16. Willey, R.L., Rutledge, R.A., Dias, S., Folks, T.,
Theodore, T., Buckler, C.E. and Martin, M.A. (1986).
Identification of conserved and divergent domains within the
5 envelope gene of the acquired immunodeficiency syndrome
retrovirus.

Proc. Natl. Acad. Sci. USA 83: 5038-5042.

10 17. Clavel, F., Guyader, M., Guétard, D., Sallé, M.,
Montagnier, L. and Alizon, M. (1986).
Molecular cloning and polymorphism of the human immune
deficiency virus type 2.

Nature 324: 691-695.

15 18. Hahn, B.H., Kong, L.I., Lee, S.-W., Kumar, P., Taylor,
M.E., Arya, S.K. and Shaw, G.M. (1987).
Relationship of HTLV-4 to simian and human immunodeficiency-
associated viruses.

Nature 300: 184-186.

20 19. Guyader, M., Emerman, M., Sonigo, P., Clavel, F.,
Montagnier, L. and Alizon, M. (1987).
Genome organization and transactivation of the human
immunodeficiency virus type 2.

25 Nature 326: 662-669.

30 20. Levy, J.A., Hoffman, A.D., Kramer, S.M., Landis, J.A.,
Shimabukuro, J.M. and Oshiro, L.S. (1984).
Isolation of lymphocytopathic retroviruses from San
Francisco patients with AIDS.

Science 225: 840-842.

35 21. Vanderborght, B., De Leys, R.J., Van Heuverswyn, H.,
Merregaert, J., Prinsen, H., Nijs, P. Vercauteren, G., and
van der Groen, G. (1988).
Submitted for publication.

-62-

1

22. Levy, J.A. and Shimabukuro, J. (1985).

5 Recovery of AIDS-associated retroviruses from patients with
AIDS or AIDS-related conditions and from clinically healthy
individuals.

J. Infect. Dis. 152: 734-738.

10

23. van der Groen, G., Vercauteren, G. and Piot, P. (1987).
Immunofluorescence tests for HIV antibody and their value as
confirmatory tests.

J. Virol. Meth. 17: 35-43.

15

24. Hoffman, A.D., Banapour, B. and Levy, J.A. (1985).
Characterization of the AIDS-associated retrovirus reverse
transcriptase and optimal conditions for its detection in
virions.

Virology 147: 326-335.

20

25. Kikukawa, R., Koyanagi, Y., Harada, S., Kobayashi, N.,
Hatanaka, M. and Yamamoto, N. (1986).
Differential susceptibility to the acquired immunodeficiency
syndrome retrovirus in cloned cells of human leukemic T-cell
line Molt-4.

(25 J. Virol. 57: 1159-1162.

30

26. Winkel, I.N., Tersmette, M., Miedema, F. and Huisman,
J.G. (1987).

Identification of gag-epitopes by a panel of MAb in a series
of HIV isolates.

Abstracts of the Third International Conference on AIDS,
Washington D.C., USA, p. 116.

35

-63-

1 27. Maizel, J.V. (1971).

Polyacrylamide gel electrophoresis of viral proteins; in
5 Methods in Virology, Vol. 5, pp: 180-246, K. Maramorosch and
H. Koprowski, Editor, Academic Press, New York, London.

28. Dunn, S.D. (1986)

Effects of the modification of transfer buffer composition
and the renaturation of proteins in gels on the recognition
10 of proteins on Western blots by monoclonal antibodies.
Anal. Biochem. 157: 144-153.

29. Gross, E. and Witkop, B. (1961).

Selective cleavage of methionyl peptide bonds in RNAase with
15 CNBr.
J. Am. Chem. Soc. 83: 1510-1511.

30. Fontana, T.P. and woods, K.R. (1981).

Modification of tryptophanyl residues with BNPS-Skatole.
20 Meth. Enzymol. 25/ 419-423.

31. Hopp, T.P. and Woods, K.R. (1981).

Prediction of protein antigenic determinants from amino acid
sequence.

25 Proc. Natl. Acad. Sci. USA 78: 3824-3828.

32. Wain-Hobson, S., Sonigo, P., Danos, O., Cole, S. and
Alizon, M. (1985).

Nucleotide sequence of the AIDS virus, LAV.
30 Cell 40: 9-17.

33. Ratner, L., Haseltine, W., Patarca, R., Livak, K.J.,
Starcich, B., Josephs, S.F., Doran, E.R., Rafalski, J.A.,
Whitehorn, E.A., Baumeister, K., Ivanoff, L., Petteway, S.R.
35 Jr., Pearson, M.L., Lautenberger, J.A., Papas, T.S.,
Ghrayes, J., Chang, N.T., Gallo, R.C. and Wong-Staal, F.

-64-

1

(1985).

Complete nucleotide sequence of the AIDS virus, HTLV-III.
Nature 313: 277-284.

5

34. Sanchez-Pescador, R., Power, M.D., Barr, P.J., Steimer, K.S., Stempien, M.M., Brown-Shimer, S.L., Gee, W.W., Renard, A., Randolph, A., Levy, J.A., Dina, D. and Luciw, P. (1985).
Nucleotide sequence and expression of an AIDS-associated
retrovirus (ARV-2)
Science 227: 484-492.

10

15

35. Muesing, M.A., Smith, D.H., Cabradillo, C.D., Benton, C.V., Laskey, L.A. and Capon, D.J. (1985).
Nucleic acid structure and expression of the human
AIDS/lymphadenopathy retrovirus.
Nature 313: 450-458.

20

36. Human retroviruses and AIDS. (1987).
G. Meyers, A.B. Rabson, S.F. Josephs, T.F. Smith and F. Wong- Staal, Editors.
Los Alamos National Laboratory, Los Alamos, New Mexico, USA.

25

37. Kawakami, T., Sherman, L., Dahlberg, J., Gazit, A., Yaniv, A., Tronick, S. and Aaronson, S.A. (1987).
Nucleotide sequence analysis of Equine Infectious Anemia
Virus proviral DNA.
Virology 158: 300-312.

30

38. Sonigo, P., Alizon, M., Staskus, K., Klatzmann, D., Cole, S., Danos, O., Retzel, E., Tiollais, P., Haase, A. and Wain- Hobson, S. (1985).

Nucleotide sequence of the Visna lentivirus: relationship to
the AIDS virus.

35

Cell 42: 369-382.

1

CLAIMS

5

10

1. HIV-3 retrovirus or variants of this virus having the essential morphological and immunological properties of any of the retroviruses deposited at the European Collection of Animal Cell Cultures (ECACC) under N° V88060301.

2. The purified retrovirus of claim 1, characterized in that said essential morphological and immunological properties are as follows:

15

20

25

30

35

- The virus exhibits a tropism for T4 lymphocytes.
- The virus is cytotoxic for the lymphocytes that it infects.
- The virus has a diameter of approximately 120 nm.
- The virus possesses a magnesium dependent reverse transcriptase activity.
- It can be cultivated in T4 receptor-bearing immortalized cell lines.
- Lysates of the virus contain a p25 protein which is immunologically distinct from the p19 protein of HTLV-I by Western blot analysis.
- Lysates of the virus contain a gp120 protein which is immunologically distinct from the gp110 protein of HTLV-I by Western blot analysis.
- The lysate of the virus contains in addition a glycoprotein with a molecular weight of 40,000 - 45,000.
- The genomic RNA of HIV-3 hybridizes neither with the sequences of HIV-1 nor with the sequences of HIV-2 under stringent hybridization conditions.

1 3. The retrovirus of claim 1 or 2, characterized in that
the nucleotide sequence of its genomic RNA which comprises
an R region and an U3 region also comprises a nucleotide
5 sequence corresponding with the following nucleotide
sequence:

	10	20	30	40	50	60
	CCCATGGATT	TGAAGATACA	CATAAAGAAA	TACTGATGIG	GAAGTTTGAT	AGATCTCTAG
	70	80	90	100	110	120
10	GCAACACCCA	TGTTGCTATG	ATAACTCAAC	CAGAGCTCCT	CCAGAAGGAC	TAAAAACTGC
	130	140	150	160	170	180
	TGAACCTGAAG	ATTGCTGACA	CTGTGGAAC	TTCAGCAAA	GACTGCTGAC	ACTGCGGGGA
	190	200	210	220	230	240
	CTTTCAGTIG	GGAGGGACAG	GGGGGGGTTT	GGGGAGTGGC	TAACCTCAG	AAGCTGCATA
15	250	260	270	280	290	300
	TAAGCAGCAG	CTTTCCTGCT	GTACGGGGTC	TOGGTTAGAG	GACCAGGTCT	GAGCCCGGGA
	310	320	330	340	350	360
	GCTCCCTGGC	CTCTAGCTGA	ACCGGCTCGT	TACGCTCAA	TAAAGCTTGC	CTTGAGTGAG

20 A.

4. The retrovirus of any of claims 1 to 3 characterized in
that its RNA virtually hybridizes neither with the Env gene
25 and the LTR close to it, in particular not with the
nucleotide sequence 8352-9538 of HIV-1, nor with the
sequences of the Pol region of the HIV-1 genome under
stringent conditions.

30 5. A composition comprising at least one antigen, in
particular a protein or glycoprotein of HIV-3 retrovirus of
any of claims 1 to 4.

6. The composition of claim 5 characterized by containing
35 a total extract or lysate of said retrovirus.

1 7. The composition of claim 5, characterized by containing
at least one of the internal core proteins of said
retrovirus, in particular p12, p16 or p26 having apparent
5 molecular weights in the order of 12,000, 16,000 and 26,000
respectively.

8. The composition of claim 5, characterized by containing
at least one of the envelope proteins of said retrovirus, in
10 particular gp41 or gp120 having apparent molecular weights
in the order of 40,000-45,000 and 120,000 respectively.

9. An antigen providing a single band in polyacrylamide gel
electrophoresis, said antigen comprising, in common with one
15 of the purified antigens of HIV-3 retrovirus, an epitope
that is recognized by serum of a patient carrying anti-HIV-3
antibodies.

10. A purified antigen having the immunological
characteristics of one of the following proteins or
20 glycoproteins of HIV-3: p12, p16, p26, gp41 and gp120.

11. The antigen of claim 10 having the aminoacid sequence,
or a part of said sequence, of the p12 protein obtained by
25 subjecting the protein mixture produced by HIV-3 to gel
electrophoresis and isolating the p12 protein in a manner
known per se.

12. The antigen of claim 10 having the aminoacid sequence,
or a part of said sequence, of the p16 protein obtained by
30 subjecting the protein mixture produced by HIV-3 to gel
electrophoresis and isolating the p16 protein in a manner
known per se.

13. The antigen of claim 10 having the aminoacid sequence,
or a part of said sequence, of the p26 protein obtained by
35 subjecting the protein mixture produced by HIV-3 to gel
electrophoresis and isolating the p26 protein in a manner
known per se.

1 14. The antigen of claim 10 having the aminoacid sequence,
or a part of said sequence, of thegp41 protein obtained by
subjecting the protein mixture produced by HIV-3 to gel
5 electrophoresis and isolating the gp41 protein in a manner
known per se.

15. The antigen of claim 10 having the aminoacid sequence,
or a part of said sequence, of thegp120 protein obtained by
10 subjecting the protein mixture produced by HIV-3 to gel
electrophoresis and isolating the gp120 protein in a manner
known per se.

16. A method for the detection of antibodies against HIV-3
15 retrovirus in a biological liquid, such as a serum or spinal
fluid, in particular for the diagnosis of a potential or
existing ARC or AIDS caused by said HIV-3 retrovirus,
characterized by contacting body fluid of a person to be
diagnosed with a composition of any of claims 5 to 8 or with
20 an antigen of any claims 9 to 15 and detecting the
immunological conjugate formed between said anti-HIV-3
antibodies and the antigen(s) used.

17. The method of claim 16, characterized in that said
25 detection of said immunological conjugate is achieved by
reacting said immunological conjugate with a labeled reagent
selected from antihuman immunoglobulin-antibodies or
bacterial A protein or G protein and detecting the complex
formed between said conjugate and said reagent.

30 18. A kit for the detection of anti-HIV-3-antibodies in a
biological fluid, comprising

- a composition as defined in any of claims 5 to 8 or an
35 antigen as defined in any of claim 9 to 15, and
- means for detecting the immunological complex formed.

1 19. The kit of claim 18, characterized in that said means
for detecting said immunological complex comprise antihuman
immunoglobulin(s) or protein A and means for detecting the
5 complex formed between the anti-HIV-3 antibodies contained
in the detected immunological conjugate.

20. An immunogenic composition containing an envelope
glycoprotein of HIV-3 retrovirus, in particular gp41 or
10 pgl20, or a part of said glycoprotein, in combination with a
pharmaceutically acceptable vehicle suitable for the
constitution of vaccines effective against HIV-3.

21. The composition of claim 20, characterized by containing
15 at least part of a glycoprotein comprising the protein
backbone of the envelope protein. or a part thereof, as
defined in any of claims 14 to 15.

22. Monoclonal antibodies characterized by their ability to
20 specifically recognize one of the antigens as defined in any
of claims 11 to 15 in particular monoclonal antibodies
specifically raised against said antigens.

23. The secreting hybridomas of the monoclonal antibodies of
25 claim 22.

24. Nucleic acids, optionally labeled, derived in part at
least of RNA of HIV-3 retrovirus or of variants thereof.

25. The nucleic acid of claim 24, characterized by
30 containing at least part of the cDNA corresponding with the
entire genomic RNA of HIV-3 retrovirus.

26. The nucleic acid of claim 24 containing the nucleotide
35 sequence as identified in claim 3.

27. The nucleic acids of claim 24 characterized by
containing nucleotide sequences coding for at least part of
the aminoacid sequences of proteins as defined in any of
claims 11 to 13.

1

28. The nucleic acids of claim 24, characterized by
containing nucleotide sequences coding for at least part of
the aminoacid sequences of glycoproteins as defined in any
of claims 14 to 15.

29. The nucleic acids of any of claims 24 to 28.
characterized by being formed into a recombinant nucleic
acid comprising a nucleic acid from a vector having said
cDNA, or a part of said cDNA, inserted therein.

30. The recombinant nucleic acid of claim 29 characterized
by being labeled.

15

31. A process for the detection of HIV-3 retrovirus or of
its RNA in a biological liquid or tissue, characterized by
contacting nucleic acids contained in said biological liquid
or tissue with a probe containing a nucleic acid according
to any of claims 25 to 30 under stringent hybridization
conditions, washing the hybrid formed with a solution
preserving said stringent conditions, and detecting the
hybrid formed.

25

32. A process for the production of HIV-3 retrovirus
characterized by culturing human T4 lymphocytes, or
permanent cell lines derived therefrom carrying the T4
phenotype, with lymphocytes or cell lines that have
previously been infected with an isolate of HIV-3
retrovirus, as well as recovering and purifying the
retrovirus from the culture medium.

30

33. A process for the production of antigens of HIV-3
retroviruses, characterized by lysing the retrovirus and
recovering the lysate containing said antigens.

35

1

34. A process for the production of any of the proteins or glycoproteins p12, p16, p26, gp41 and gp120 as defined hereinbefore, or of a part thereof, characterized by inserting the corresponding nucleic acid sequence in an expression vector, transforming a host with said vector, culturing the transformed host as well as recovering and purifying the expressed protein.

5

10

35. A process for the production of a hybridization probe for the detection of the RNA of HIV-3 retrovirus, characterized by inserting a DNA sequence, particularly of any of claims 24 to 29 in a cloning vector by in vitro recombination, cloning the modified vector obtained in a suitable cellular host, and recovering the hybridization probe.

15

20

36. A method for detecting antigen of HIV-3, characterized by coating a surface with an immunoglobulin fraction raised against HIV-3, bringing a body or culture fluid to be analyzed into contact with the immunoglobulins, and detecting the complex formed between the immunoglobulins and the antigen.

25

30

35




Fig. 1

SUBSTITUTE SHEET




Fig. 2

SUBSTITUTE SHEET




Fig. 3

SUBSTITUTE SHEET

4/19




Fig. 4

SUBSTITUTE SHEET




Fig. 5

SUBSTITUTE SHEET

6/19





Fig. 6


SUBSTITUTE SHEET

7/19



SUBSTITUTE SHEET

Fig. 7



SUBSTITUTE SHEET

Fig. 8




Fig. 9

SUBSTITUTE SHEET





Fig. 10

SUBSTITUTE SHEET

11/19

WELLS COATED WITH:

- HIV-1 lysate
- HIV-3 (ANT 70) lysate
- ▲—▲ HIV-2 lysate

Fig. 11

SUBSTITUTE SHEET

12/19

A. p17 gag




Fig. 12a

SUBSTITUTE SHEET

B. p24 gag





Fig. 12b

SUBSTITUTE SHEET

C. pol



— conserved protease sequence
 ^ cleavage site protease/reverse transcriptase
 | cleavage site reverse transcriptase/endonuclease

Fig. 12c

SEQUENCE SHEET





Fig. 13a



SUBSTITUTE SHEET

16/19






Fig. 13b

SUBSTITUTE SHEET




markers

Fig. 13c



SUBSTITUTE SHEET

p53/65 reverse transcriptase




BNPS-Skatole cleavage



Fig. 13d

markers

p31 endonuclease



SUBSTITUTE SHEET

A. STRINGENTB. NONSTRINGENTVIRUS SPOTTED:

A ANT 70 RNA
 B HIV-1(SF 4) RNA
 C HIV-2 (53) RNA

ML. EQUIVALENT CULTURE SUPERNATANT:

1. 5 ml.
2. 2.5 ml.
3. 1.3 ml.
4. 0.6 ml.

Fig. 14

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No PCT/EP 89/00643

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC ⁴ C 12 N 7/00, A 61 K 39/21, G 01 N 33/569, C 12 N 15/00, IPC: A 61 K 39/395, C 12 Q 1/68, C 12 Q 1/70		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁴	C 12 N, A 61 K, G 01 N	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP, A, 0239425 (INSTITUT PASTEUR) 30 September 1987 cited in the application --	
A	WO, A, 86/02383 (INSTITUT PASTEUR) 24 April 1986 cited in the application --	
A	AIDS Research and Human Retroviruses, vol. 23, no. 1, 1987, Mary Ann Liebert, Inc., Publishers, G. Franchini et al.: "Genetic analysis of a new subgroup of human and simian T-lymphotropic retroviruses: HTLV-IV, LAV-2, SBL-6669, and STLV-III AGM", pages 11-17 --	
A	Science, vol. 233, 18 July 1986, F. Clavel et al.: "Isolation of a new human retrovirus from West African patients with AIDS", pages 343-346 cited in the application -- ./.	
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
1st September 1989	02. 10. 89	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	T.K. WILLIS	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
------------	--	----------------------

- | | | |
|---|--|--|
| A | Nature, vol. 324, no. 18, 25 December 1986,
F. Clavel et al.: "Molecular cloning and polymorphism of the human immune deficiency virus type 2", pages 691-695
cited in the application | |
| A | Nature, vol. 326, 16 April 1987,
M. Guyader et al.: "Genome organization and transactivation of the human immunodeficiency virus type 2", pages 662-669
cited in the application | |
| A | Nature, vol. 330, 12 November 1987,
B.H. Hahn et al.: "Relation of HTLV-4 to simian and human immunodeficiency-associated viruses", pages 184-186 | |
| A | Nature, vol. 333, no. 6172, 2 June 1988,
M. Fukasawa et al.: "Sequence of simian immunodeficiency virus from African green monkey, a new member of the HIV/SIV group", pages 457-461 | |

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

EP 8900643
SA 29055

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 22/09/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0239425	30-09-87	FR-A- 2593189	24-07-87
		FR-A- 2593190	24-07-87
		FR-A,B 2593922	07-08-87
		FR-A,B 2594229	14-08-87
		FR-A,B 2596063	25-09-87
		FR-A,B 2597500	23-10-87
		AU-A- 6891187	14-08-87
		EP-A- 0320495	14-06-89
		WO-A- 8704459	30-07-87
		JP-T- 63502242	01-09-88
		OA-A- 8468	29-07-88
		US-A- 4839288	13-06-89
		EP-A- 0269520	01-06-88
WO-A- 8602383	24-04-86	FR-A,B 2571968	25-04-86
		AU-A- 5061785	02-05-86
		EP-A- 0201540	20-11-86
		JP-T- 62500592	12-03-87
		AU-A- 5320086	13-08-86
		WO-A- 8604336	31-07-86
		EP-A- 0211022	25-02-87
		JP-T- 62502095	20-08-87
		OA-A- 8413	30-06-88

EPO FORM 10479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)